Jul 15 357

Mr. Jeff Luedke Green Light Company P.O. Box 17985 San Antonio, TX 78217

Dear Mr. Luedke:

Subject: Resubmission

Green Light Dursban Granules EPA Registration No. 869-158

Your Application Dated April 21, 1987

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable; however, you must delete the statement "This product contains the toxic ingredient methylene chloride" since you are not increasing the amount of inert ingredients.

Submit five (5) copies of draft labels, deleting the aforementioned statement.

Sincerely yours,

Dennis H. Edwards, Jr. Product Manager (12) Insecticide-Rodenticide Branch Registration Division (TS-767C)

	85156: I	/C:Edwards +	E-10.KENCO:	6/9/87:6/1	8/87:ron:vo	ek:ron		
SYMBOL	,							
SURNAME		 						
DATE								
EPA Form	1320-1 (4-81)	<u> </u>	L	l	L	I	OFF	CIAL FILE CORY

GREEN LIGHT DURSBANR GRANULES

FOR USE ON LAWNS, IN FLOWER BEDS, AND UNDER SHRUBBERY

For Control Of Fleas, Ticks, Ants, Chinch Bugs, Brown Dog Ticks, Crickets, Sod Webworms, Cutworms, Sowbugs, Pillbugs, Earwigs, Hyperodes (turfgrass) Weevils

15 Lb. Bag Granules Treats 3,000 Square Feet

ACTIVE INGREDIENT:

 Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)

 phosphorothioate]
 0.5%

 INERT INGREDIENTS
 99.5%

 Total
 100.0%

Contains 3/16-1b. Chlorpyrifos per gallon Dursban^R registered trademark The Dow Chemical Company

THIS PRODUCT CONTAINS THE TOXIC INGREDIENT METHYLENE CHLORIDE

is and delde

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ COMPLETE PRECAUTIONS ON BACK PANEL

EPA REG. NO. 869-158

EPA EST. NO. 869-TX-1

With COMMENTS in HPA Letter Dated:

Formulated by Green Light Co., San Antonio, TX 78217

Net Weight 15 LBS.

JUN 15:987

i Inspettoide

anticide Ag

ie posticide

To Fig. No.

Under the

Pur Las

as . .

remsia:

(Back Panel)

CAUTION

KEEP OUT OF REACH OF CHILDREN

May by harmful if swallowed. Do not take internally. Avoid eye contact. Flust contaminated eyes with plenty of water, and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse.

PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote.

USE PRECAUTIONS: Keep children and pets off treated areas until material is washed into soil and grass is dry. Keep away from food, feedstuffs, and domestic warer supplies.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes: Apply this product only as specified on this label.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. PESTICIDE DISPOSAL: Leave unwanted pesticide material in original container. Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard in trash.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

Apply Green Light Durstan^R Granules insecticide uniformly to home lawns, flower beds, under shrubbery, and other ornamental and recreational turf grass areas using hand fertilizer spreader or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using Granules at the rate of 5 pounds per 1,000 sq.ft. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application sprinkle the turf with 150 gallons of water per 3,000 sq. ft. to wash the granules into the zone of insect infestation. Retreat as necessary.

NOTE: See directions on fertilizer or granule spreader for setting to apply 5 pounds per 1,000 square feet.

RATES TO USE

Area to be Treated	Pounds of Granules	Gallons of Water to Sprinkle on Turf after Treatment			
1,000 sq. ft.	5	50			
3,000 sq. ft.	15	150			

To control Hyperodes (turfgrass) weevils use twice the rate of Granules shown on table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.