

GREEN LIGHT MANY PURPOSE DUST

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

KILLS ANTS - Very Effective for control of Ants, Digger Wasps, Chinch Bugs, Armyworms, Clover and Bermuda Mites, Crickets, Cutworms, Earwigs, Billbugs, Sod Webworms, Aphids, Leafhoppers, Cucumber Beetles, Leafminers, Flea Beetles, Mealybugs, Whiteflies, Thrips, Tent Caterpillars, Wireworms, Mole Crickets, Fleas, Ticks, Chiggers

ACTIVE INGREDIENT:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....

5%

INERT INGREDIENTS.....

95%


Total

100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Additional Cautions On Back



EPA Reg. No. 869-159

EPA Est. No. 869-TX-1

Green Light Company San Antonio, TX 78127

Net Weight 1 lb.

(Back Panel)

READ ENTIRE LABEL BEFORE USING

USE THIS PRODUCT ONLY AS DIRECTED

CAUTION: Harmful if swallowed. Avoid breathing dust. May be absorbed through skin. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL CAUTION: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or to residues. Obtain protective information from your Cooperative Agricultural Extension Service. Keep children and pets off treated area until this material is washed into the soil and grass is dry.

PHYSICIAN: Toxic symptoms which may be observed after accidental ingestion: headache, giddiness, blurred vision, nervousness, weakness, cramps, diarrhea, discomfort in the chest, sweating, miosis, tearing, salivation, and other excessive respiratory tract secretion, vomiting, cyanosis, papilledema, uncontrollable muscle twitches, convulsions, coma, loss of sphincter control.

Antidote: Atropine is antidotal providing it is used as soon as early poisoning symptoms appear and the dose of Diazinon is not too great. It is important to give large doses of atropine, up to ten times the usual dosage.

DIRECTIONS FOR USE

Ants, Digger Wasps: Dust directly onto ant hills and into wasp ground nest openings.

PESTS IN YARD:

Spread amount indicated evenly over 100 sq. ft. of lawn. Ants, Armyworms, Clover Mites, Crickets, Cutworms, Earwigs, Billbugs, Webworms, use 4 ounces (8 tablespoons). Bermuda Mites, Chiggers, Fleas, Ticks, Leafhoppers, use 1½ ounces (2½ tablespoons). Chinch Bugs in St. Augustine lawns, use 7 ounces (14 tablespoons).

GARDEN INSECTS:

General Directions - When insects or their damage first appear apply this product to plants so that a thin film of dust covers both upper and under sides of foliage. Repeat applications as needed to maintain effective control.

For flowers, shrubs, and ornamental gardens - aphids, flea beetles, leafhoppers, leafminers, mealybugs, mites, whiteflies, scale crawlers, thrips, tent caterpillars, bagworms. NOTE: Do not use on ferns, poinsettias, hibiscus, papaya, pilea, gardenia, since injury may occur.

VEGETABLE GARDENS:

NOTE: Do not use on food crops within the number of days of harvest shown in parentheses after each crop: Tomatoes (3), Potatoes (14): aphids, banded cucumber beetles, armyworms, leafminers. Lettuce (10), Radishes (10), Spinach (10), Turnips (10), Peas (0), Beans (7): aphids, leafhoppers, Mexican bean and cucumber beetles, mites.

SOIL INSECTS:

Cutworms and wireworms on lima and snap beans, peas, peppers, tomatoes, collards, and in flower gardens - Before planting spread 1 lb. of Dust evenly over 800 sq. ft. of area to be planted, or 2 oz. (4 tablespoons) per 100 sq. ft. immediately after spreading, mix thoroughly into soil to depth indicated: cutworms, 2-3 inches; subterranean cutworms, 3-6 inches; wireworms, 4-8 inches.

Mole Crickets on broccoli, cabbage, cauliflower, collards, endive, lettuce, peppers, radishes, strawberries, tomatoes, turnips. One or two days before seeding or transplanting, spread 4 oz. (8 tablespoons) evenly over 500 sq. ft. of area to be planted and mix into top 1 or 2 inches of soil.

NOTICE: Seller makes no warranty of any kind, express or implied, concerning effects of use of this product other than those specified on this label. Buyer, or user, accepts all responsibility for results due to misuse of this product.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. PESTICIDE DISPOSAL: Unwanted pesticide material leave in original container. Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse container. Discard in trash.