

GREEN LIGHT
BIO-WORM KILLER
contains *Thuricide-

FOR HOME GARDENS: Biological Insecticide for control of certain leaf-chewing worms on Vegetables, Fruits, Shade Trees and Ornamentals listed on label.

ACTIVE INGREDIENTS:

<u>Bacillus thuringiensis</u> Berliner potency of 4,000 international units (at least 6 million viable spores) per milligram**	0.8%
Petroleum Hydrocarbon Solvent	3.0%
INERT INGREDIENTS	96.2%
Total	100.0%

*Thuricide- Registered trademark of Sandoz Inc.
**Equivalent to 4.0 billion International Units per quart.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK FOR ADDITIONAL PRECAUTIONS

EPA Est. No. 869-TX-1

EPA Reg. No. 869-174

869-174

Distributed by Green Light Co., San Antonio, TX 78217

Net Contents 1 Pt.
8 Oz.

(Back or Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes, nose, mouth. Avoid breathing spray mist.

ENVIRONMENTAL HAZARDS: Keep out of lake, pond, or stream. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. Activity may be impaired at temperatures above 90 F. Store in a cool place. Protect from freezing. PESTICIDE DISPOSAL: Empty pesticide container into a container of new paper and discard in trash. CONTAINER DISPOSAL: Do not reuse container. Recycle the container before re-use in the home.

SHAKE OR STIR BEFORE USING -- Pour recommended amount of Bio-Worm Killer into a partially filled sprayer. Stir and add the remaining amount of water. Shake sprayer occasionally during spraying. Do not allow diluted spray to stand in sprayer for more than 12 hours. Combinations with chemical insecticides, fungicides or spreader-stickers in the sprayer are generally not harmful to Bio-Worm Killer if used properly. Good under and upper leaf coverage is essential since the insect must ingest Bio-Worm Killer. Bio-Worm Killer must remain on leaf surfaces to be effective. Re-apply after heavy rains. Susceptible insects stop feeding after eating Bio-Worm Killer, but may otherwise appear to be unaffected for several days.

VEGETABLES and FRUITS: Mix 1.5 fluid ounces of Bio-Worm Killer per 3 gallons of water (1 tablespoon per gallon) for a hand sprayer and apply to all plant foliage to cover an area of 1,000 sq. ft. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following insects:

TOMATO HORNWORM on tomatoes.

CABBAGE LOOPER on broccoli, cauliflower, collards, kale, mustard greens, turnip green, cabbage, celery, lettuce, melons and tomatoes.

IMPORTED CABBAGE WORM on broccoli, cabbage, cauliflower, collards, kale, mustard greens and turnip greens.

GRAPE LEAF FOLDER and OMBIVOROUS LEAF ROLLER (California Only) on Grapes.

FRUIT TREE LEAFROLLER and ORANGE DOG (California Only) on Oranges. SHADE

TREES and ORNAMENTALS: Mix 2 fluid ounces of Bio-Worm Killer per 3 gallons of water (4 teaspoons per gallon) for a hand sprayer or a proportional amount for a hose-end sprayer following manufacturer's directions.

Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms:

Spring Caterpillar, Fall Caterpillar, Fall Webworm, Oak Moth Larva (California only), Red-humped Caterpillar (California Only) and Tent Caterpillar.