


869-174

12/4/2001

Page 1 of 3



GREEN LIGHT
BT WORM KILLER

BIOLOGICAL INSECTICIDE FOR CONTROL OF CERTAIN LEAF-CHEWING WORMS ON VEGETABLES, FRUITS, SHADE TREES AND ORNAMENTALS

ACTIVE INGREDIENT:

Bacillus thuringiensis variety *kurstaki* Potency of 4,000 International Units) At least 6 million viable spores per mg*)..... 0.8%

OTHER INGREDIENTS..... 99.2%

Total 100.0%

*Equivalent to 4.0 billion International Units per quart

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back or Side Panel For Additional Precautionary Statements

EPA Reg. No. 869-174

EPA Est. No. 73273-TX-1

Manufactured by Southwest Contract Packaging for Green Light Sales Company
 P.O. Box 17985, San Antonio, TX 78217-0985 www.greenlightco.com

NET CONTENTS ONE PINT (474 ml)

(Back or Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes or open wounds. Avoid breathing spray mist.

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> . Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> . Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

PRODUCT INFORMATION

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Green Light Company at (210) 494-3481 for general information and/or medical information.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water when disposing of washwater rinsate.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. Activity may be impaired at temperatures above 90 F. Store in a cool place. Protect from freezing.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap in several layers of newspaper and discard in trash.

GENERAL INFORMATION: SHAKE WELL BEFORE USING. Pour recommended amount of Green Light BT Worm Killer into a partially filled sprayer. Shake and add the remaining amount of water. Shake sprayer occasionally during spraying. Do not allow diluted spray to stand in sprayer for more than 12 hours. Good under and upper leaf coverage is essential since the worm must ingest Green Light BT Worm Killer. Susceptible worms stop feeding after eating Green Light BT Worm Killer, but may otherwise appear to be unaffected for several days. Re-apply after heavy rains. Best results are obtained by treatments when worms are small; they must be actively feeding on treated, exposed foliage. You may apply Green Light BT Worm Killer up to and on the day of harvest. Kills worm and caterpillar stage insects, but has no effect on humans, pets, birds, earthworms or beneficial insects such as honeybees and ladybugs.

VEGETABLES AND FRUITS: Mix 1 tablespoon of Green Light BT Worm Killer in 1 gallon of water and apply to all plant foliage. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following pests:

TOMATO HORNWORM on tomatoes.

CABBAGE LOOPER on broccoli, cauliflower, collards, kale, mustard greens, turnip greens, cabbage, celery, lettuce, melons and tomatoes.

IMPORTED CABBAGE WORM on broccoli, cabbage, cauliflower, collards, kale, mustard greens and turnip greens.

OMNIVOROUS LEAF ROLLER on Grapes.

FRUIT TREE LEAFROLLER and ORANGE DOG on Oranges.

SHADE TREES AND ORNAMENTALS: Mix 4 teaspoons in 1 gallon of water in a hand sprayer or a hose-end sprayer. Apply at the first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms: Spring Cankerworm, Fall Cankerworm, Fall Webworm, Bagworm and Tent Caterpillar.