

JUN 16 1989

Mr. Jeff Luedke
Green Light Company
P.O. Box 17985
San Antonio, TX 78217

Dear Mr. Luedke:

Subject: Green Light Rotenone
EPA Registration No. 869-186
Your Submission of a Draft Label With a Cover Letter Dated
September 8, 1989 Responding to the Registration Standard
for Rotenone

Enclosed is a stamped copy of the draft label you submitted. It will be acceptable with the following change. Revise the "Pesticide Disposal" statement to read "Leave unused pesticide in the original container. Securely wrap the original container in several layers of newspaper and discard in trash."

Within three (3) months of the date stamped on this letter, submit a revised Confidential Statement of Formula which includes certified limits.

Sincerely yours,



Phil Hutton
Product Manager (18)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

50560:I:Schroeder:C.Disk:KEVRIC:06/17/91:aw:wo:aw

CONCURRENCES

SYMBOL	SURNAME	DATE					

293

GREEN LIGHT ROTENONE

Kills Insects On Vegetables, Fruits, And Ornamentals. Also For Control Of Fleas On Dogs And Cats.

ACTIVE INGREDIENTS:

Rotenone.....	0.75Z
Other Associated Resins.....	1.50Z
INERT INGREDIENTS.....	97.75Z
Total	100.00Z

JUL 16 1991

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-186

EPA Est. No. 869-TX-1

Formulated by Green Light Co., San Antonio, TX 78217

Net Wt. 1 lb.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Do not get in eyes or on skin. Wash thoroughly after handling. Avoid contamination of food and feed. Do not use on humans.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED, call a physician immediately. IF INHALED, remove victim to fresh air. IF ON SKIN, wash with soap and water. IF IN EYES, rinse with plenty of water. Call a physician.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. **PESTICIDE DISPOSAL:** Unwanted pesticide material leave in original container. Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse container. Discard in trash.

For Use On Vegetables, Fruits, And Ornamentals

Apply this product when infestation first appears. Apply at a rate of 1/2 to 2/3 lbs. per 1,000 sq. ft. Do not apply within one day of harvest. Reapply as necessary.

VEGETABLES:

Asparagus---Asparagus Beetle, Japanese Beetle
Beans (Snap & Lima)---Aphids, Bean Leaf Beetle, Cucumber Beetle, Flea Beetle, Mexican Bean Beetle

BEST AVAILABLE COPY

375
Beets---Flea Beetle

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards---Cabbage Looper, Diamond Back Moth Larvae, Flea Beetle, Imported Cabbageworm

Cantaloupe, Cucumbers, Squash, Watermelon---Cucumber Beetle, Melon Worm, Plant Lice
Eggplant---Aphids, Colorado Potato Beetle, Flea Beetle

Lettuce---Loopers

Peas---Pea Aphid, Pea Weevil

Potatoes---Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetle

Radishes---Flea Beetle

Spinach---Flea Beetle, Spinach Aphid, Spinach Lice

Tomato---Aphid, Colorado Potato Beetle, Flea Beetle, Tomato Hornworm, Tomato Suck Fly

Turnips---Flea Beetle

FRUITS:

Apples, Peaches---Japanese Beetle

Blackberries, Blueberries, Dewberries, Grapes, Raspberries---Japanese Beetle, Raspberry Fruit Fly, Raspberry Fruit Worm, Sawfly. For Raspberry Cane Borer apply to foliage just before plants bloom and repeat in 7 to 10 day intervals.

Currants, Gooseberries---Gooseberry Fruit Worm

Strawberries---Flea Beetle, Leaf Beetle, Spittlebug

ORNAMENTALS: Apply thoroughly as needed for control of the following insects:

Aphids---Aster, Gladiolus, Iris, Rose, Sunflower, Sweet Peas,

Cabbage Looper And Imported Cabbage Worm---Carnation, Chrysanthemum, Dahlia, Daisy, Rose, Sweet Peas, Zinnia

Flea Beetle---Petunia, Primrose, Sunflower, Violet, Wallflower

Japanese Beetle---Rose

Fleas On Dogs And Cats: Dust animal thoroughly and rub into fur. Repeat weekly as required for control. Do not allow product to get into eyes, nose or ears of animals. Do not use on puppies or kittens less than 4 weeks old.

Fleas In Kennels, Pens, Lawns, And Under Houses: Apply 5 to 10 ounces per 250 sq. ft. Remove debris before dusting.

Lice On Horses: Apply 1 to 4 ounces of powder depending on the size of the animal. Work into hair. Repeat in 10 to 12 days and as needed thereafter.

BEST AVAILABLE COPY