

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

**CAUTION**

May cause irritation of nose, throat, eyes and skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Wash thoroughly after using and before eating or smoking. Remove and wash contaminated clothing before reuse.

Do not contaminate water, food or feed by storage or disposal.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL AND CHEMICAL HAZARDS**

Keep away from fire and sparks.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Mix 1 level tablespoon per 1 gallon of water. Mix with 3 parts oil to get best results. Do not use oil if you are not sure.

**STORAGE AND DISPOSAL**

**STORAGE:** Keep out of reach of children and pets. Store in a cool place, but not below 32°F. Do not allow to become overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this product. Keep container closed when not in use. Leave in original container until used.

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

**TOMATOES, GREENHOUSE VEGETABLES:** Late blight, Early blight, Septoria leaf spot, Late blight, Early blight, Septoria leaf spot, Late blight, Early blight, Septoria leaf spot. Apply when first signs appear. Repeat at 7 to 10 day intervals for 3 or 4 applications or until disease is controlled. If control is not achieved, more frequent spraying may be necessary. If necessary, spray before harvest. Do not spray within 8 days of harvest.

**POTATOES:** Early blight, Late blight. Spray 9 to 12 days throughout the season. Begin spray when plants are 2 to 3 inches high.

**BEANS:** Anthracnose, Downy mildew, Rust. Start spraying when disease first appears and repeat at 7 to 10 day intervals. Do not spray on succulent beans within 8 days of harvest.

**CUCUMBERS:** Anthracnose, Downy mildew, Leaf Spot, Angular leaf spot, Ascochyta blight, Gummy stem blight, Phytophthora blight. Begin when disease first appears. Repeat at 7 to 10 day intervals. Do not spray within 8 days of harvest.

**PEPPERS:** Anthracnose, Downy mildew, Leaf Spot, Phytophthora blight, Rust. Begin when disease first appears. Repeat at 7 to 10 day intervals throughout the season.

**CUCUMBERS:** Downy mildew, Leaf Spot, Anthracnose, Phytophthora blight. Begin when disease first appears. Repeat at 7 to 10 day intervals.

**EGGPLANT:** Anthracnose, Early blight, Late blight, Phytophthora blight. Begin when 2 to 3 inches high. Repeat at 7 to 10 day intervals throughout the season.

**NECTARINES & PEACHES:** Brown rot, Shot hole. Begin when first signs appear. Repeat at 7 to 10 day intervals. Do not spray within 14 days of harvest.

**CARROTS:** Downy mildew, Rust. Start spraying when plants are 8 to 12 inches high. Repeat after harvest and again 10 days later. Do not spray within 7 days of harvest.

**DESSERTS:** Spray when plants are well rooted out and apply at 7 to 10 day intervals throughout the season.

**BEANS:** Downy mildew, Leaf Spot, Anthracnose, Phytophthora blight. Begin when plants are well rooted out and repeat as needed. Do not spray within 8 days of harvest.

**CUCUMBERS:** Downy mildew, Leaf Spot, Anthracnose, Phytophthora blight. Begin when plants are well rooted out and repeat as needed. Do not spray within 8 days of harvest.

**PEPPERS:** Downy mildew, Leaf Spot, Anthracnose, Phytophthora blight. Begin when plants are well rooted out and repeat as needed. Do not spray within 8 days of harvest.

**TOMATOES:** Downy mildew, Leaf Spot, Anthracnose, Phytophthora blight. Begin when plants are well rooted out and repeat as needed. Do not spray within 8 days of harvest.

**POTATOES:** Downy mildew, Late blight. Spray 9 to 12 days throughout the season. Begin spray when plants are 2 to 3 inches high.



**GREEN LIGHT  
LIQUID FLOWABLE  
FUNGICIDE**

**CONTROLS MANY DISEASES OF FRUITS, VEGETABLES,  
LAWNS AND ORNAMENTALS**

**ACTIVE INGREDIENT:**  
Maneb (Manganese ethylene bis(dithiocarbamate) 37.0%  
(Total Manganese) 7.6%  
**INERT INGREDIENTS** 63.0%  
Total 100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panel for additional precautionary statements.

E.P.A. Reg. No. 869-190

E.P.A. Est. No. 869-TX-1

Distributed by GREEN LIGHT COMPANY, San Antonio, Texas 78217

NET CONTENTS 1 PINT

ACCEPTED  
with COMMENTS  
to EPA Letter Patent

NOV 24 1982

Under the Fungicide...  
Registered...  
869-190