

869-215

11/7/2003

1/1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

If inhaled - Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. - Call a poison control center or doctor for further treatment advice.

If on skin or clothing - Take off contaminated clothing. - Rinse skin with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.

If in eyes - Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.

If swallowed - Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by poison control center or doctor. - Do not give anything by mouth to an unconscious person.

PRODUCT INFORMATION

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for general information and/or medical information.

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION: The active ingredient, *Bacillus thuringiensis*, is a naturally occurring component of the environment. Green Light Dipel Dust is a highly selective insecticide for use against listed caterpillars (larvae) of Lepidopterous insects. Honey bees foraging treated areas are not harmed when Dipel Dust is used according to label directions.

APPLICATION DIRECTIONS: Since Dipel Dust works only in a caterpillar's stomach, a small amount of treated leaf must be eaten to provide a lethal dose. Complete coverage of both sides of foliage is necessary. After biting a treated portion of the leaf, caterpillars stop feeding within a few hours. Death follows in a few days.

Apply when caterpillars (or their damage) first appear. Retreatment at 7 or 10 day intervals may be required, especially as egg hatch may occur over an extended period of time. Use the higher rates against heavy infestations, or where it may be difficult to obtain good coverage. Dipel Dust may be applied up to time of harvest.

APPLICATION RATES

CROP	PEST	SZ./AQ FT. ROW
Lettuce, Broccoli, Cabbage, Cauliflower, Collards, Spinach, Turnip Greens, Cerey, Beans, Peas	Loopers	1 1/4-2 1/2
	Hornworm	3/4-1 1/4
Tomatoes, Peppers, Eggplant	Imported Cabbageworm	3/4-1 1/4
	Moth	1-2
Green	Green	1-2
	Cloverworm	3/4-1 1/4
	Webworm	1 1/4-2 1/2
	Amyworms*	2 1/2-5

*Dipel Dust may be used to suppress small armyworms (1st & 2nd instar) when populations are light and full ground coverage can be attained.

Tomatoes, Peppers, Eggplant	Loopers	1 1/4-2 1/2
	Hornworm	3/4-1 1/4
Tomato	Fruitworm*	2 1/2
	Variegated Cutworm	2 1/2
Caterpillars	Salt Marsh	1 1/4-2 1/2
	Amyworms**	2 1/2-5

*Apply weekly in a preventative program. **Dipel Dust may be used to suppress small armyworms (1st & 2nd instar) when populations are light and full ground coverage can be attained.

Cucumbers, Melons, Cucumbers, etc.	Ringworm	1 1/4-2 1/2
	Complex* Melonworms	2 1/2-5

*Apply weekly in a preventative program.

Livestock	Soil Webworm	4-8 ozs/100 sq. ft.
-----------	--------------	---------------------

Flowers and Ornamentals	Loopers	1 1/4-2 1/2
	Tobacco Budworm	1 1/4-2 1/2

Omnivorous	Loopers	1 1/4-2 1/2
	Leafroller	1 1/4-2 1/2

Omnivorous	Loopers	1 1/4-2 1/2
	Leafroller	1 1/4-2 1/2

Moth	Armyworms*	1-1 1/4
	Elk Moth (Hornworm)	2 1/2-5

Lo Moth	Lo Moth	3/4-1 1/4
	Oleander Moth	1 1/4-2 1/2


*Dipel Dust may be used to suppress small armyworms (1st & 2nd instar) when populations are light and full coverage ground sprays are applied

EPA Reg. No. 869-215 EPA Est. No. 73273-TX-1

Manufactured For
GREEN LIGHT COMPANY
P.O. Box 17983
San Antonio, TX 78217-0985
www.greenlightco.com Tel (210) 494-3481



Rev. 09-23-2003



Contains *Bacillus thuringiensis* (B.T.)
Biological Insecticide For Control of Tomato
Hornworm, Looper, Webworm, Armyworm,
Cutworm, Leaf Roller, Tomato Fruitworm
and Other Caterpillars On Vegetables,
Fruits, Ornamentals, Flowers,
and Lawns

ACTIVE INGREDIENT:

<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> solids, spores and lipidolipran active toxins	0.49%
OTHER INGREDIENTS:	99.51%
TOTAL	100.00%

The percent active ingredient does not indicate product performance and potency rates are not federally standardized.

DIPEL® is a registered trademark of Abbott Laboratories

KEEP OUT OF THE REACH OF CHILDREN.
CAUTION See Back Panel for Additional Precautionary Statements

NET WT. 1 LB. (454gm)

