

Reg # 869-216

PM-25

1 of 4



GREEN LIGHT COMPANY

P.O. BOX 17985

SAN ANTONIO, TEXAS 78217-0985

January 29, 1993

Joanne I. Miller, Product Manager 23
U.S. Environmental Protection Agency
Office of Pesticide Programs
Registration Division (H7505C)
401 M St., S.W.
Washington, DC 20460

RE: CERTIFICATION STATEMENTS AND REVISED LABELING FOR GREEN LIGHT
WIPE OUT, EPA REG. NO. 869-2164

Dear Ms. Miller:

I, being an authorized representative of Green Light Company, certify that all containers of Green Light Wipe Out, EPA Reg. No. 869-216 produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Sincerely,

GREEN LIGHT COMPANY

A handwritten signature in black ink, appearing to read "Jeff Luedke".

Jeff Luedke
Director of Regulatory Affairs
and Product Development

enclosure

GREEN LIGHT WIPE-OUT

FOR BROADLEAF WEED CONTROL IN LAWNS AND TURF GRASSES

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichloro- phenoxyacetic Acid*.....	3.26%
Dimethylamine Salt of 2-(2-Methyl-4-Chloro- phenoxy) Propionic Acid**.....	3.28%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy) Propionic Acid***.....	3.24%

INERT INGREDIENTS..... 90.22%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid.....	2.72%
**2-(2-Methyl-4-Chlorophenoxy)Propionic Acid..	2.72%
***2-(2,4-Dichlorophenoxy) Propionic Acid.....	2.72%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-216 EPA Est. No. 869-TX-1
 Net Contents One Quart (0.946 liters)
 Green Light Co., San Antonio, TX 78217-0985

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin.

STATEMENT OF PRACTICAL TREATMENT: IF IN EYES: Immediately flush eyes with plenty of water for 15 minutes and get medical attention immediately. IF ON SKIN: Wash promptly with soap and water and rinse thoroughly. IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger.

ENVIRONMENTAL HAZARDS: Avoid drift of spray onto vegetables, flowers, shrubs, trees or other desirable plants. This product is toxic to fish and aquatic invertebrates. Drift or run-off may adversely affect fish and aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment wash water.

PROTECTIVE CLOTHING: When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wear face shield or goggles when mixing or applying this product. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

RE-ENTRY STATEMENT: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

STORAGE: Keep from freezing. Store in a locked storage area out of reach of children and domestic animals. Store in original container. **PESTICIDE DISPOSAL:** Leave unwanted pesticide material in original container. Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

GENERAL INFORMATION: Apply spray when weeds are small and tender for best results. For older and hard-to-kill weeds use higher dilution rate. When St. Augustine is growing rapidly or in cases of accidental overdose, some temporary discoloration of the foliage may occur. **DO NOT USE ON DICHONDRA OR NEWLY SEEDED LAWNS NOR ON LAWNS WHERE DESIRABLE CLOVERS ARE PRESENT.** Newly seeded lawns may be treated after they have been mowed at least three times and are well established. Spray when air temperature is between 50-90 F. Avoid applying during hot or dry periods unless irrigation is used a few days before.

Do not water for 24 hours after treatment. Spray works faster with temperature above 70 F. Spray when air is calm to avoid drift. Spray will injure clover, dichondra, most ornamental and vegetable crops. This product works slowly. Effects begin to show after a few days and weeds gradually die. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in injury.

HOW TO MIX: (1) For use on St. Augustine grass, Bentgrass, Buffalograss and grass seedlings, mix 1 - 1/2 ounce (3 tablespoons) in 1 gallon of water and uniformly spray 300 square feet. (Example: an area 10 feet wide and 30 feet long). If necessary, repeat treatment in 2 to 3 weeks, if hard-to-kill weeds persist. (2) For use on all other grasses such as Bermuda, Bahia, Bluegrass, Centipede, Fescue, Rye, mix 2 ounces (4 tablespoons) in 1 gallon of water and spray uniformly over 300 square feet. On hard-to-kill species, or under poor growing conditions, or on older weeds, mix 2 - 1/2 ounces in 1 gallon of water and spray on 300 square feet with special attention to thorough coverage of hard-to-kill weeds. The maximum number of broadcast applications per treatment site is 2 per year.

WEEDS CONTROLLED: Black Medic, Buckhorn, Burdock, Buttercup, Canada Thistle, Carpetweeds, Chickweed, Chicory, Clover,

Cocklebur, Dandelion, Dock, Dog Fennel, English Daisy, Florida Pursley, Frenchweed, Ground Ivy, Hawkweed, Heal-All, Heartloaf Drymary, Henbit, Jimsonweed, Knotweed, Lambsquarter, Mallow, Morningglory, Mustard, Oxalis, Pennywort, Pepperweed, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Red Clover, Red Sorrel, Sheep Sorrel, Smartweed, Speedwell, Spurge, Spurweed, Stitchwort, Toadflax, Veronica, Wild Aster, Wild Carrot, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Onion, Wild Radish, Wood Sorrel, Yarrow, and many other broadleaf weeds.

1950
1951
1952
1953
1954