

Reg # 869-222

PM-22

Jacket

1977



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 M St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
869-222

Date of Issuance:
FEB 18 1994

Term of Issuance:
Conditional

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

Name of Pesticide Product:
Green Light Fung-Away
Systemic Fungicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Green Light Company
P.O. Box 17985
San Antonio, TX 78217-0985

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Add the phrase "EPA Registration No. 869-222" to your label before you release the product for shipment.
3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Signature of Approving Official:

[Handwritten signature]

Date:

[Handwritten date]

297

page 2
EPA Reg. No. 869-222

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

) Enclosure

)

397
Jacket Copy

GREEN LIGHT FUNG-AWAY®

Systemic Fungicide

Contains Bayleton® Fungicide

FOR CONTROL OF POWDERY MILDEW, LEAF BLIGHT/SPOTS, RUSTS, ANTHRACNOSE, BROWN PATCH AND OTHER DISEASES ON FLOWERS, SHRUBS, TREES AND LAWNS

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone..... 0.88%

INERT INGREDIENTS..... 99.12%
100.00%

Contains 0.15 ounces by weight of 1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone per pint
U.S. Patent No. 3,912,752

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

FEB 18 1984

Bayleton® is a Registered Trademark of the ultimate parent company of Miles Inc.

Fung-Away® is a Registered Trademark of Green Light Company

Under the Federal Insecticide, Fungicide Act and the pesticide laws of the state of California
EPA Reg. No.

869-222

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-

EPA Est. No. 869-TX-1

Formulated by Green Light Co., P.O. Box 17985 San Antonio, TX 78217-0985

Net Contents One Pint (474 ml)

(Back or Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled. Avoid breathing vapor or spray mist. This product may cause allergic skin reactions. Keep out of reach of children. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT: IF EYES ARE CONTAMINATED, flush eyes with plenty of water. Call a physician if irritation persists. IF ON SKIN, wash skin immediately with soap and warm water. If irritation occurs, get medical attention. IF INHALED, remove to fresh air. If not breathing, give artificial respiration preferably mouth to mouth. Get medical attention. IF SWALLOWED, administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

BEST AVAILABLE COPY

IN CASE OF POISONING, call a physician or Poison Control Center. Have patient lie down and keep quiet.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

SYMPTOMS OF POISONING: No specific symptoms. Acute poisoning is accompanied by hyperactivity followed by sedation.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not make applications when weather conditions favor drift from target area.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION: Green Light Fung Away is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage is necessary. Rainfall or sprinkler irrigation, within 1/2 hour after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply in all cases when plant root systems are established and actively growing. Applications should be applied at recommended intervals to maintain disease control.



Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or lawns.

RECOMMENDED APPLICATIONS TURFGRASS DISEASE CONTROL

DISEASES	OUNCES PER GALLON PER 1,000 SQ. FT.		REMARKS
	Preventative	Curative	
Dollar Spot	7		PREVENTATIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of FUNG AWAY may extend for as long as 60 days dependent upon environmental conditions.
		14	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventative schedule and rate.
Copper Spot	7		PREVENTATIVE RATE: Apply at 15 to

Powdery Mildew
 Red Thread
 (Laetisaria
fuciformis)
 Rusts

30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of FUNG AWAY can be greater than 30 days depending on environmental conditions.

Brown Patch
 (Rhizoctonia
solani) 14
 (Suppression)

CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventative schedule and rate.

Anthracnose 14 14
 Poa Annua
 Decline

Apply at 30 day intervals as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.

PREVENTATIVE RATES ONLY

Stripe Smut 14

Make a single application in Spring as grass growth begins.

Fusarium Blight 14
 (Fusarium
culmorum)
 (Fusarium
poae)
 Summer Patch
 (Phialophora
graminicola)

Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.

ORNAMENTAL PLANT DISEASE CONTROL

PLANTS (Disease Controlled Reference)

FLOWERING AND FOLIAGE PLANTS (Outdoor)	SHADE TREES	DISEASES
Ageratum (2b,3,4)	Ash (3)	(1) Flower Blight
Aster (4)	Aspen (3,4)	a. <u>Ovulinia</u> spp.*
Eegonia (3)	Balsam (4)	b. <u>Sclerotinia</u> spp.*
Canna (4)	Birch (3,4)	c. <u>Colletotrichum</u> *
Carnation (3,4)	Buckeye (3)	(2) Leaf Blight/Spots
Chrysanthemum (3,4)	Chestnut (3)	a. <u>Cephalosporium</u> spp.***
Dahlia (3)	Cottonwood (3,4)	b. <u>Cercospora</u> spp.
Delphinium (3)	Elm (3)	c. <u>Didymella</u> spp.**
Dianthus (4)	Fir (4)	d. <u>Entomosporium</u>
Four O' Clock (4)	Locust (3)	
Geranium (3,4)	Maple (3)	
Hollyhock (3,4)	Paulownia (3)	

- Iris (2c)
- Marigold (2b,4)
- Nephtytis (2a)
- Pansy (3,4)
- Petunia (3,4)
- Phlox (2b,3,4)
- Rose (3)
- Salvia (3,4)
- Sedum (3)
- Snapdragon (3,4)
- Sunflowers (3,4)
(ornamentals only)
- Zinnia (2b,3)

- Oak (3)
- Pine (4,5)
- Poplar (3,4)
- Russian Olive (2b,4)
- Sycamore (3)
- Vitex (2b)
- Walnut (3)
- Willow (3,4)

- Hawaii Only
- Dendrobium (1c)

spp.***

- (3) Powdery Mildew
 - a. Erysiphe spp.
 - b. Microsphaera spp.
 - c. Oidium spp.
 - d. Podosphaera spp.
 - e. Phyllactinia spp.
 - f. Sphaerotheca spp.
 - g. Uncinula spp.

FLOWERING FOLIAGE AND WOODY SHRUBS

- Amelanchier (3)
- Azalea (1a,3)
- Barberry (3,4)
- Buckthorn (4)
- Camellia (suppression of 1b)
- Crabapple (Flowering) (3,4)
- Crape Myrtle (3)
- Dogwood (3)
- Euonymus (3)
- Gardenia (3)
- Hawthorn (3,4)
- Holly (3)
- Juniper (4)
- Leucothoe (2b)
- Lilac (3)
- Mock Orange (3,4)
- Mountain Laurel (1a,2b,3)
- Nineoark (3)
- Pear (Flowering) (3)
- Photinia (2d,3,4)
- Potentilla (4)
(Cinquefoil)
- Privet (2b,3)
- Pyracantha (3)
- Rhododendron (1a,2b,3)
- Rose (3)
- Spirea (3)
- Viburnum (3,4)

- (4) Rusts
 - a. Coleosporium spp.
 - b. Cronartium spp.**
(Fusiform)
 - c. Gymnosporangium spp.
 - d. Melampsora spp.
 - e. Melampsoridium spp.
 - f. Peridermium spp.** (Gall)
 - g. Phragmidium andersonii
 - h. Puccinia spp.
 - i. Uromyces spp.
 - j. Uredinopsis mirabilis*

- (5) Tip Blight
 - a. Sirococcus strobilinus**

APPLICATION RATES; Except as noted for specific diseases, mix 1 to 2 tablespoonfuls (1/2 to 1 fl. oz.) of Fung-Away in one gallon of water, and apply as a full coverage foliage spray to the point of drip as needed.

*Mix 2 to 4 fl. ozs. of Fung-Away in one gallon of water and apply as a full coverage foliar spray to the point of drip. Applications for flower blight

should begin at the expanded bud stage (color showing). Use multiple applications at 7 to 14 day intervals as needed dependent upon bloom periods.

**Mix 4 fl. ozs. of Fung-Away plus sufficient spreader sticker for good coverage in one gallon of water. Apply in a spray application to the point of run-off on an as needed basis. Excessive rates or excessive applications may result in a shortening of the flower stalk on iris.

***Mix 2 to 4 fl. ozs. of Fung-Away in one gallon of water and apply as a full coverage foliar spray to point of run-off. Apply in early spring as growth starts and reapply on a 14 to 21 day interval until new growth is fully expanded. Protect new growth that develops in late summer or fall as temperatures begin to drop. ✓

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.

