

PM 22

869-224

page 1 Jacket

GREEN LIGHT FUNG-AWAY® SYSTEMIC LAWN FUNGICIDE

FOR CONTROL OF DISEASES ON TURF

Contains BAYLETON® Fungicide

1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended for the pesticide registered under EPA Reg. No. 869-224

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone..... 0.5%

INERT INGREDIENTS..... 99.5%
100.0%

Bayleton® is a Registered Trademark of the ultimate parent company of Miles Inc.

Fung-Away® is a Registered Trademark of Green Light Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-224

EPA Est. No. 869-TX-1

Formulated by Green Light Co., P.O. Box 17985 San Antonio, TX 78217-0985

Net Weight 8 lbs. (3.6 kg)

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION; Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED, administer water freely and induce vomiting by giving one dose (½ or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately. IF ON SKIN, wash skin immediately with soap and warm water. If irritation occurs, get medical attention. IF IN EYES ARE CONTAMINATED, wash with plenty of water for at least 15 minutes. Get medical attention. IF INHALED, remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically. In case of poisoning, notify Green Light Company (512)494-3481.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do

BEST AVAILABLE COPY

not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION: Green Light Fung-Away Lawn Fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control. Do not use on crops grown for food or forage.

RECOMMENDED APPLICATIONS TURFGRASS DISEASE CONTROL

Disease	Lbs. of Fung-Away per 1,000 sq. ft. Preventative Curative	Remarks
Dollar Spot	3	Preventative Rate: Apply recommended rate at 30 day intervals. Protective activity may extend as long as 60 days dependent upon environmental conditions.
	6	Curative Rate: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to preventative rate schedule for rates and timing.
Copper Spot Powdery Mildew Red Thread (Laetisara fuciformis) Rusts	3	Preventative Rate: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Fung-Away can be greater than 30 days depending on environmental conditions.
Brown Patch (Rhizoctonia solani) (Suppression)	6	Curative Rate: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedule for rates and timing.

BEST AVAILABLE COPY

Anthracnose 3
Poa Annua
Decline

Preventative Rate: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.

6 Curative Rate: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedule for rates and timing.

Take-All Patch
(Gaeumannomyces)
graminis var.
graminis)
(Except California)

Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.

DM
3125-413

3 Preventative Rate: Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 to 28 day intervals.

6 Curative Rate: To control existing infections, apply 6 lbs. at 14 to 21 day intervals for the initial 1 or 2 treatments followed by 3 lbs. at 21 to 28 day intervals. Cultural control practices such as aerification, top-dressing, reseeding, and fertilization must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.

Preventative Rates Only

Stripe Smut	3	Make a single application in Spring as grass growth begins.
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer Patch (Phialophora graminicola) (except California)	3	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer Patch (Phialophora graminicola) (California only)	3 to 6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 3 lb. rate on 15 day intervals or at the 6 lb. rate on 30 day intervals.
Gray Snow Mold (Typhula incarnata)(Except California)	6	Apply in Fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold Fusarium Patch (Fusarium nivale) Southern Blight (Sclerotium rolfsii) (Except California)	3 to 6	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover or when turf is dormant. Use higher rate in areas with a history of severe disease damage

SPREADER SETTINGS AND APPLICATION RATES

IMPORTANT: THESE SETTINGS ARE ONLY APPROXIMATE. AGE AND CONDITION OF SPREADER CAN CAUSE WIDE VARIATION.

For best results, cover the area 2 times, in cross directions, at 1/2 rate. This method ensures complete coverage, as the second covering provides an opportunity for correction of possible operational errors or dial setting errors made on the first covering.

Spread widths given on chart are operational widths based on brisk walking. Maintain an EVEN, NORMAL pace. A faster or slower pace, or tipping spreader forward or backward, will vary the spread width.

MAKE	MODEL	LBS. PER 1,000 SQ. FT.			SPREAD WIDTH
		1	3	6	
Broadcast Spreaders					
CYCLONE	B1/B55	3	4-3/4	5-1/2	6'
ORTHO	6000	1	3	7	6'
SCOTTS	ER-2	24-3/4	27-1/2	29-1/2	3-1/2'
SEARS 1100	671.1987	2	3-3/4	6	6'
Drop Spreaders					
ORTHO	5000	4-1/4	5-3/4	7-3/4	18"
SCOTTS	PF-3	5	6	8	21"

NOTE: Make applications when turf is dry and following the mowing operation. Do not mow until a minimum of 1/4 inch of water has been applied to treated area. Do not use clippings for animal feed.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **HOUSEHOLD:** Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area.