



869-235

3-25-2003

1/7

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

259579

Application for Pesticide - Section I

1. Company/Product Number 869-235	2. EPA Product Manager LaRocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Green Light Conquest Insecticide Concentrate	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Green Light Company P.O. Box 17985 San Antonio, TX 78217-0985 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

MAR 25 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Revised Ready-To-Spray directions. Please see highlighted text on attached label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willingly make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)				
Name Nora M. Hodge		Title Registration Specialist		
		Telephone No. (Include Area Code) (210) 494-3481		
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)	
2. Signature 		3. Title Registration Specialist		
4. Typed Name Nora M. Hodge		5. Date 3-12-2003		

2/7

Fast-Acting Water-Based Long-Lasting Low Odor Non-Staining

Provides Up To One Month (4 Weeks) Control Of Named Insect Pests Outdoors

(OPTION B) READY-TO-SPRAY FOR CONVENIENT AND ACCURATE APPLICATION. CONTROLS NAMED INSECT PESTS OUTSIDE AROUND HOMES, ON HOME LAWNS, TREES, SHRUBS AND FLOWERS.

Permethrin	[*(3-Phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate].....	2.5%
------------	--	------

***cis/trans isomer ratio:** Min. 35% (+) cis
Max. 65% (+) trans

Conquest is a registered trademark of Green Light Company

CAUTION

See Side or Back Panel For Additional Precautionary Statements

Manufactured for Green Light Company, P.O. Box 17985, San Antonio, TX 78217-0985
www.greenlightco.com (210) 494-3481

EPA Reg. No. 869-235

EPA Est. No. 73273-TX-1

Net Contents One Quart (0.946 liters)

NOTIFICATION

MAR 25 2003

3/7

(Back or Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse.

FIRST AID

If inhaled

- . Move person to fresh air.
- . If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- . Call a poison control center or doctor for further treatment advice.

If on skin or clothing

- . Take off contaminated clothing.
- . Rinse skin immediately with plenty of water for 15-20 minutes.
- . Call a poison control center or doctor for treatment advice.

If in eyes

- . Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- . Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- . Call a poison control center or doctor for treatment advice.

If swallowed

- . Call poison control center or doctor immediately for treatment advice.
- . Have person sip a glass of water if able to swallow.
- . Do not induce vomiting unless told to do so by the poison control center or doctor.
- . Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for general information and/or medical information.

4/7

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. Do not contaminate ornamental fish ponds. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **SHAKE WELL BEFORE USING.** Not for use on plants grown for sale or other commercial use, or for commercial feed production or for research purposes. Not for use in greenhouses, sod farms or golf courses.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. **DISPOSAL: If empty-**Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled-**Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

VEGETABLES

Celery: For control of Armyworms (Beet, Fall, Southern), Cabbage Loopers, Cutworms (Black, Granulate) and Leafminers, mix 1-1/2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply every 3 to 5 days or as needed. May be applied up to 3 days of harvest. Do not make more than 10 applications per season.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower: For control of Cabbage Aphids, Beet Armyworms, Cabbage Loopers, Diamondback Moths and Imported Cabbageworms, mix 1 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5 days or as needed. Can be applied up to 1 day of harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli, and no more than 10 applications on Cabbage.

Eggplant: For control of Colorado Potato Beetles and Leafminers, mix 3 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Do not exceed 16 applications per season. Can be applied up to 3 days of harvest.

Head Lettuce: For control of Corn Earworms, Beet Armyworms, Loopers (Alfalfa, Cabbage) and Tobacco Budworms, mix 2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5 to

10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day of harvest.

Potatoes: For control of Beetles (Colorado Potato, Potato Flea), Beet Armyworms, Cabbage Loopers, Cutworms, European Corn Borers, Leafhoppers (Aster, Potato), Potato Aphids, Potato Psyllids, Potato Tuberworms and Tarnished Plant Bugs, mix 1-1/2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed. Do not make more than 12 applications per season. Can be applied up to 7 days of harvest.

Peppers: For control of Cabbage Loopers, Corn Earworms, European Corn Borers, Flea Beetles, Pepper Weevils and Leafminers, mix 2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days of harvest.

Spinach: For control of Beet Armyworms, Cabbage Loopers, Corn Earworms, Cutworms, European Corn Borers, Green Cloverworms and Leafhoppers, mix 1-1/2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply when needed, but not more than 7 applications per season. Can be applied up to 7 days of harvest.

Tomatoes: For control of Tomato Fruitworms, Cabbage Loopers, Tomato Pinworms, Armyworms (Beet, Southern), Colorado Potato Beetles, Granulate Cutworms and Tomato Hornworms, mix 1-1/2 fl. ozs. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed up to day of harvest. Do not make more than 6 applications per season. Do not use on tomato varieties in which the mature tomatoes will be less than 1 inch in diameter (such as cherry tomatoes).

FRUITS AND NUTS

General Directions: Mix 2 fl. ozs. in 1 gallon of water. Spray when insect pests first appear. Wet trees to the dripping point. Try to make application to undersides of leaves and penetrate dense foliage. Make spray applications in the early morning or late afternoon on days when the winds are light.

Almonds: For control of Navel Orangeworms and Peach Twig Borers. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days of harvest.

Apples: For control of Green Fruitworms, Leafrollers (Oblique-banded, Redbanded), Plum Curculio, Rosy Apple Aphids, Spotted Tentiform Leafminers, Tarnished Plant Bugs and White Apple Leafhoppers. Do not make more than 3 applications per season. Do not apply after petal fall.

Peaches: For control of Borers (Lesser Peachtree, Peach Twig), Green Fruitworms, Plum Curculio, Oriental Fruit Moths and Tarnished Plant Bugs. Do not make more than 8 applications per season. Can be applied up to 7 days of harvest.

Pears: For control of Pear Psylla. Apply during dormant through delayed dormant growth period only. Do not make more than 2 applications per season. **Pears (summer):** For control of Codling Moths, Green Fruitworms and Pear Psylla. Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

TREES AND SHRUBS

General Directions: Mix 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Make spray application to the underside of leaves and try to

6/7

penetrate dense foliage. Spray at the first sign of insect pest infestation. Make applications in the early morning or late afternoon on days when wind is light. Repeat treatments as necessary.

For use on a variety of trees and shrubs including: Azalea, Birch, Elm, Euonymus, Fir, Honeysuckle, Lilac, Oak, Palm, Pine, Poplar. For control of insect pests including: Aphids, Bagworms, Budworms, Cicadas, Elm Leaf Beetles, Exposed Thrips, Fall Cankerworms, Fall Webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Oakworms, Pine Beetles, Pine Moths, Scale, Spider Mites, Whiteflies.

ROSES AND FLOWERS

General Directions: Mix 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting the plants to the dripping point. Make application to the underside of leaves and try to penetrate dense foliage. Make applications in the early morning or late afternoon on days when wind is light. Spray at the first sign of insect pest infestation. Repeat treatments as necessary.

For use on a variety of flowers including: Ageratum, Aster, Azalea, Begonia, Coleus, Gladiolus, Marigold, Orchid, Pansy, Petunia, Rose, Snapdragon, Zinnia. For control of insect pests including: Aphids, Bagworms, Cicadas, Exposed Thrips, Fall Webworms, Inchworms, Leafhoppers, Leafminers, Mealybugs, Spider Mites, Whiteflies.

As with all plants listed above, especially chrysanthemums and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

HOME LAWNS

For use on Bent, Bluegrass, Bermuda, Dichondra, Fescue, St. Augustine, to control Ants (including foraging Fire Ants*), Armyworms, Chiggers, Chinch Bugs, Crickets (including Mole Crickets), Cutworms, Fleas, Grasshoppers, Sod Webworms (Lawn Moths), Ticks, White Grubs of: Japanese Beetle, European Chafer and Southern Chafer. Thoroughly wet down grass a few hours before applying. Mix 6 fl. ozs. in 10 gallons of water to treat 1,000 sq. ft. of home lawn area. For White Grubs, water lawn thoroughly after application. Treat anytime between late July and early October. For ants, spot spray ant mounds.

NOTE: Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

*Aids in Control

(OPTION B)

READY-TO-SPRAY DIRECTIONS FOR OUTDOOR USE ONLY ON HOME LAWNS:

The sprayer attached to the container is ready-to-use. Simply attach to your garden hose and follow these simple instructions:

(1) Insure the large, round, "on/off" knob is set to OFF.

- (2) Turn on water.
- (3) Rotate and PUSH IN small knob near front of sprayer. This allows product to mix with water.
- (4) Hold sprayer and container level and point towards area to be sprayed.
- (5) Turn large, round, "on/off" knob forward (away from you) to ON.
- (6) Begin spraying product evenly over area you wish to treat.
- (7) To stop spraying, turn large, round, "on/off" knob backward (toward you) to OFF.
- (8) PULL OUT and rotate small knob near front of sprayer. This prevents product from mixing with water.
- (9) Turn off water at the faucet. Relieve water pressure in the hose by turning large, round, "on/off" knob to ON until water pressure is reduced.
- (10) Turn the knob to OFF for storage of unused product and/or disposal of empty container.

HOME LAWNS

For use on turfgrass varieties including: Bent, Bluegrass, Bermuda, Dichondra, Fescue, St. Augustine, to control Ants (including foraging Fire Ants*), Armyworms, Chinch Bugs, Sod Webworms (Lawn Moths), Fleas, Ticks, Crickets, Cutworms, Earwigs, Grasshoppers, Chiggers, Mosquitoes and Pillbugs (Sowbugs) infesting home lawns. Wet grass down thoroughly before spraying. Spray when insect pests first appear. Retreat as needed. Uniformly apply 6 fl. ozs. per 1,000 sq. ft. of home lawn area. Refer to fill level markings on side panel of container to determine amount (ounces) of product used. Do not allow use of treated areas by pets or people during application or until spray has dried. DO NOT USE ON PETS.

*Aids in Control

ROSES, FLOWERS, TREES, SHRUBS

For control of Aphids, Bagworms, Tent Caterpillars, Fall Webworms, Grasshoppers, Spider Mites, Whiteflies, Exposed Thrips, Mealybugs, Japanese Beetles, Leafminers, Leafrollers, Scale. Avoid over-spraying to the point of excessive runoff. Treat when insect pests first appear and retreat as needed.

NUISANCE INSECT PESTS OUTSIDE THE HOME

For control of Ants (including Fire*, Pharaoh (Sugar), Carpenter, Red Harvester, Odorous House), Clover Mites, Crickets, Earwigs, Millipedes, Pillbugs (Sowbugs), Fleas, Ticks, Flies, Gnats, Mosquitoes, Spiders, Scorpions, Chiggers, Roaches. Spray window and door screens, porches, patios, entranceways, walks, driveways, and into cracks and other places where insect pests can hide. To prevent entry into homes, spray a 5 foot band of soil around the home next to the foundation and spray the foundation of the home to a height of 2 to 3 feet. Repeat as needed.

*Aids in Control

