

869-235

8/27/2008

10f7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Jeff Luedke
Green Light Company
P.O. Box 17985
San Antonio, TX 78217-0985

AUG 27 2008

Dear Mr. Luedke:

Subject: Amendment –Add lawn use precaution, and
add Ready-to-Spray Directions for Outdoor Use Only
Green Light Conquest Insecticide Concentrate
EPA Reg. No. 869-235
Your submission dated April 10, 2008

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) section 3(c)(7)(a), is acceptable and a stamped copy of the label is enclosed for your records.

Sincerely,

George LaRocca.
Product Manager 13
Insecticide Branch
Registration Division (7505P)

2087

GREEN LIGHT CONQUEST INSECTICIDE CONCENTRATE

FOR USE ON RESIDENTIAL LAWNS AND GARDENS ONLY

Fast-Acting Water-Based Long-Lasting Low Odor Non-Staining

ONE QUART COVERS OVER 5,000 SQ. FT. OF HOME LAWN AREA

Provides Up To One Month (4 Weeks) Control Of Named Insect Pests Outdoors

FOR USE ON HOME LAWNS, TREES, SHRUBS, ROSES,
FLOWERS, FRUITS, NUTS AND VEGETABLES

(OPTION B) READY-TO-SPRAY FOR CONVENIENT AND ACCURATE APPLICATION. CONTROLS NAMED INSECT PESTS OUTSIDE AROUND HOMES, ON HOME LAWNS, TREES, SHRUBS AND FLOWERS.

ACTIVE INGREDIENT:

Permethrin [* (3-Phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate]..... 2.5%

OTHER INGREDIENTS..... 97.5%
TOTAL 100.0%

*cis/trans isomer ratio: Min. 35% (±) cis
Max. 65% (±) trans

Conquest® is a registered trademark of Green Light Company.

KEEP OUT OF REACH OF CHILDREN

CAUTION


See Booklet For Additional Precautionary Statements

Manufactured for:
Green Light Company
P.O. Box 17985
San Antonio, TX 78217-0985

www.greenlightco.com
Tel. (210) 494-3481

EPA Reg. No. 869-235
EPA Est. No. 73273-TX-1

Net Contents One Quart (0.946 liters)



3067

(BOOKLET)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse.

FIRST AID

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a Poison Control Center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for general information and/or medical information.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. Do not contaminate ornamental fish ponds. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

Not for use on plants grown for sale or other commercial use, or for commercial feed production, or for research purposes. Not for use in greenhouses, sod farms, or golf courses.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. **DISPOSAL: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

VEGETABLES

Celery: For control of Armyworms (Beet, Fall, Southern), Cabbage Loopers, Cutworms (Black, Granulate), and Leafminers, mix 1½ fl.oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 3 to 5 days, or as needed. May be applied up to 3 days of harvest. Do not make more than 10 applications per season.

Broccoli, Brussels Sprouts, Cabbage, and Cauliflower: For control of Cabbage Aphids, Beet Armyworms, Cabbage Loopers, Diamondback Moths, and Imported Cabbageworms, mix 1 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5 days or as needed. Can be applied up to 1 day of harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli, and no more than 10 applications on Cabbage.

Eggplant: For control of Colorado Potato Beetles and Leafminers, mix 3 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Do not exceed 16 applications per season. Can be applied up to 3 days of harvest.

Head Lettuce: For control of Corn Earworms, Beet Armyworms, Loopers (Alfalfa, Cabbage), and Tobacco Budworms, mix 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply every 5 to 10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day of harvest.

Potatoes: For control of Beetles (Colorado Potato, Potato Flea), Beet Armyworms, Cabbage Loopers, Cutworms, European Corn Borers, Leafhoppers (Aster, Potato), Potato Aphids, Potato Psyllids, Potato Tubeworms, and Tarnished Plant Bugs, mix 1½ fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed. Do not make more than 12 applications per season. Can be applied up to 7 days of harvest.

Peppers: For control of Cabbage Loopers, Corn Earworms, European Corn Borers, Flea Beetles, Pepper Weevils, and Leafminers, mix 2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days of harvest.

Spinach: For control of Beet Armyworms, Cabbage Loopers, Corn Earworms, Cutworms, European Corn Borers, Green Cloverworms, and Leafhoppers, mix 1½ fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed, but not more than 7 applications per season. Can be applied up to 7 days of harvest.

Tomatoes: For control of Tomato Fruitworms, Cabbage Loopers, Tomato Pinworms, Armyworms (Beet, Southern), Colorado Potato Beetles, Granulate Cutworms, and Tomato Hornworms, mix 1½ fl. oz. in 1 gallon of water to cover 1,000 sq. ft. Apply as needed up to day of harvest. Do not make more than 6 applications per season. Do not use on tomato varieties in which that mature tomatoes will be less than 1 inch in diameter (such as cherry tomatoes).

FRUITS AND NUTS

General Directions: Mix 2 fl. oz. in 1 gallon of water. Spray when insect pests first appear. Wet trees to the dripping point. Try to make application to undersides of leaves and penetrate dense

foliage. Make applications in the early morning or late afternoon on days when the winds are light.

Almonds: For control of Navel Orangeworms and Peach Twig Borers. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days of harvest.

Apples: For control of Green Fruitworms, Leafrollers (Oblique-banded, Redbanded), Plum Curculio, Rosy Apple Aphids, Spotted Tentiform Leafminers, Tarnished Plant Bugs, and White Apple Leafhoppers. Do not make more than 3 applications preseason. Do not apply after petal fall.

Peaches: For control of Borers (Lesser Peachtree, Peach Twig), Green Fruitworm, Plum Curculio, Oriental Fruit Moths, and Tarnished Plant Bugs. Do not make more than 8 applications per season. Can be applied up to 7 days of harvest.

Pears: For control of Pear Psylla. Apply during dormant through delayed dormant growth period only. Do not make more than 2 applications per season. **Pears (Summer):** For control of Codling Moths, Green Fruitworms, and Pear Psylla. Do not make more than 3 applications per season. Can be applied up to 14 days of harvest.

TREES AND SHRUBS

General Directions: Mix 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Make spray application to the underside of leaves and try to penetrate dense foliage. Spray at the first sign of insect pest infestation. Make applications in the early morning or later afternoon on days when wind is light. Repeat treatments as necessary.

For use on a variety of trees and shrubs including: Azalea, Birch, Elm, Euonymus, Fir, Honeysuckle, Lilac, Oak, Palm, Pine, and Poplar. For control of insect pests including: Aphids, Bagworms, Budworms, Cicadas, Elm Leaf Beetles, Exposed Thrips, Fall Cankerworms, Fall Webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Oakworms, Pine Beetles, Pine Months, Scale, Spider Moths, and Whiteflies.

ROSES AND FLOWERS

General Directions: Mix 1 fl. oz. in 1 gallon of water. Apply as a thorough spray, wetting the plants to the dripping point. Make application to the underside of leaves and try to penetrate dense foliage. Make applications in the early morning or late afternoon on days when wind is light. Spray at the first sign of insect pest infestation. Repeat treatments as necessary.

For use on a variety of flowers including: Ageratum, Aster, Azalea, Begonia, Coleus, Gladiolus, Marigold, Orchid, Pansy, Petunia, Rose, Snapdragon, and Zinnia. For control of insect pests including: Aphids, Bagworms, Cicadas, Exposed Thrips, Fall Webworms, Inchworms, Leafhoppers, Leafminers, Mealybugs, Spider Mites, and Whiteflies.

As with all plants listed above, especially Chrysanthemums and Roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

6007

HOME LAWNS

For use on Bent, Bluegrass, Bermuda, Dichondra, Fescue, St. Augustine, to control Ants (including foraging Fire Ants*), Armyworms, Chiggers, Chinch Bugs, Crickets (including Mole Crickets), Cutworms, Fleas, Grasshoppers, Sod Webworms (Lawn Moths), Ticks, White Grubs of: Japanese Beetle, European Chafer, and Southern Chafer. Thoroughly wet down grass a few hours before applying. Mix 6 fl. oz. in 10 gallons of water to treat 1,000 sq. ft. of home lawn area. For White Grubs, water lawn thoroughly after application. Treat anytime between late July and early October. For Ants, spot spray ant mounds.

*Aids in control.

NOTE: Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

(OPTION B)


READY-TO-SPRAY DIRECTIONS FOR USE OUTSIDE AROUND HOMES, ON HOME LAWNS, TREES, SHRUBS, ROSES AND FLOWERS

The sprayer attached to the container is ready-to-use. Simply attach to your garden hose and follow these simple instructions:

READY-TO-SPRAY DIRECTIONS FOR OUTDOOR USE ONLY.

The sprayer attached to the container is ready-to-use. Simply attach to the garden hose and follow these instructions:

1. Turn on water.
2. Hold sprayer and container level and point towards area to be sprayed.
3. Bend yellow tab back and turn knob backward (toward you) to ON.
4. Begin spraying product evenly over area you wish to treat.
5. To stop spraying, turn knob forward (away from you) to OFF.
6. Turn off water.
7. Properly store unused product or dispose of empty container.



ROSES, FLOWERS, TREES, SHRUBS

For control of Aphids, Bagworms, Tent Caterpillars, Fall Webworms, Grasshoppers, Spider Mites, Whiteflies, Exposed Thrips, Mealybugs, Japanese Beetles, Leafminers, Leafrollers, Scale. Avoid over-spraying to the point of excessive runoff. Treat when insect pests first appear and retreat as needed.

NUISANCE INSECT PESTS OUTSIDE THE HOME

For control of Ants (including Fire*, Pharaoh (Sugar), Carpenter, Red Harvester, Odorous House), Clover Mites, Crickets, Earwigs, Millipedes, Pillbugs (Sowbugs), Fleas, Ticks, Flies,

707

Gnats, Mosquitoes, Spiders, Scorpions, Chiggers, Roaches. Spray window and door screens, porches, patios, entranceways, walks, driveways, and into cracks and other places where insect pests can hide. To prevent entry into homes, spray a 5 foot band of soil around the home next to the foundation and spray the foundation of the home to a height of 2 to 3 feet. Repeat as needed.

*Aids in Control