

BACK PANEL

DIRECTIONS FOR USE
(Read Carefully)

DILUTION RATE: The general rate of application is 1 lb. in 20 gallons of water (3 tablespoons per gallon). Note heavier rates required for bagworm and scale insect control.

WHEN TO APPLY: For most effective results follow a regular spray program in early Spring when insects first appear, but before diseases appear. Continue at 10 day intervals, after heavy rains, or as needed. Cover both upper and lower surfaces of foliage. For timing of application on fruit trees follow suggested schedule or consult your State Agricultural Extension Service for additional information consistent with label directions and cautions.

RECOMMENDED USES ON ROSES, ORNAMENTALS, SHRUBS AND TREES:

DISEASES - For the control of Black Spot and Powdery Mildew of Roses; Septoria Leaf Spot on Chrysanthemums; Rust of Carnations; Powdery Mildew and Anthraxnose of Zinnias, Marigolds, Phlox, Snapdragons and Asters.

INSECTS - For the control of Aphids, Spider Mites, Japanese Beetles, Thrips, Lacebugs, Leaf Rollers, Whitefly, Leafminers on Boxwood, Holly and Birch; Tent Caterpillars, Cankerworms.

When used for the control of Bagworms on Spruce, Euonymous Scale Crawlers, Soft Brown Scale Crawlers, Azalea Scale Crawlers on Azalea, Rhododendron, and other shrubs use 1 lb. in 10 gallons of water (6 heaping tablespoons per gallon).

NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula.
Do not use on Petunias or Boston Ivy.
Do not use on Japanese Maples.

RECOMMENDED USES ON FRUIT AND VEGETABLES:

DISEASES - For the control of Apple Scab, Sooty Blotch, Bitter Rot, Leaf Spot on Apples, Crabapples, and Sour Cherries; Downy and Powdery Mildew on Cucumbers and Melons; Fruit Rot on Blackberries and Raspberries, Anthraxnose and Grapes.

INSECTS - For the control of Codling Moth, Curculio, Spider Mites, Pear Psylla and Apple Maggot on APPLES; Tent Caterpillars, Black Cherry Aphid, Japanese Beetles and Two Spotted Mites on CHERRIES.

Japanese Beetles, Grape Berry Moth, Grape Leafhopper and Red Banded Leaf Roller on GRAPES.

Japanese Beetle, Leaf Rollers, Leafhoppers and Raspberry Aphids on BLACKBERRIES and RASPBERRIES.

Aphids, Spider Mites, Flea Beetles and Colorado Potato Beetles on TOMATOES.
Aphids, Cucumber Beetles and Squash Bugs on CUCUMBERS, MELONS and SQUASH.

CAUTION: DO NOT APPLY TO VEGETABLES WITHIN 2 DAYS OF HARVEST - 5 DAYS ON TOMATOES, 7 DAYS ON BLACKBERRIES AND RASPBERRIES.

SUGGESTED FRUIT SPRAY SCHEDULE

Delayed Dormant.....(From first green on bud to 1/2 inch leaves on Apple)

Pre-Bloom.....(From first pink to first bloom on Apple and Cherry)

Petal Fall.....(When last petals have fallen) on all fruits listed above.

Cover Sprays.....(2 to 4 applications) begin 10 days after petal fall on Apples.

Post Harvest Spray..For Cherry Leaf Spot apply immediately after harvest and 10 days later.

CAUTIONS: DO NOT APPLY TO FRUIT TREES WITHIN 7 DAYS OF HARVEST AND ON SOUR CHERRIES MAKE 1 APPLICATION AND THEN NOT AGAIN FOR 30 DAYS.
MAY BE HARMFUL TO SOME VARIETIES OF APPLES.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Harmful if swallowed. Do not breathe dust. Do not take internally. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear long sleeved work clothing. Change to clean clothing daily. Note for physicians: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

Do not contaminate or store near feed or foodstuffs. This product is toxic to fish. Keep out of any body of water.

DISCLAIMER

The seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of these products are beyond the control of the seller. Bayer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.