

875-47

11-26-2007

1/10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460



November 26, 2007

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Kresti A. Lyddon  
Senior Registration Specialist  
JohnsonDiversey, Inc.  
8310 16<sup>th</sup> Street, MS 486  
Sturtevant, WI 53177-0902

Subject: Dibac  
EPA Registration No. 875-47  
Application Date: October 24, 2007  
Receipt Date: November 6, 2007

Dear Ms. Lyddon:

This acknowledges receipt of your notification, submitted under the provision of PR Notice 98-10, FIFRA section 3(c)9.

**Proposed Notification**

- Addition of National Sanitation Foundation logo

**General Comments**

Based on a review of the material submitted, the following comment applies:

The notification application is acceptable and a copy has been inserted in your file for future reference.

Should you have any questions or comments concerning this letter, please contact me at (703) 308-6345.

Sincerely,

Wanda Y. Henson  
Product Reviewer (32)  
Regulatory Management Branch II  
Antimicrobials Division (7510P)

CONCURRENCES

SYMBOL	7510P	7510P					
URNAM	E. Berg	Henson					
DATE	11/26/07						

2/10

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060 Approval Expires 5-31-98



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number <b>875-47</b>	2. EPA Product Manager <b>Emily Mitchell</b>	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>Dibac™</b>	PM# <b>32</b>	
5. Name and Address of Applicant (Include ZIP Code) <b>JohnsonDiversey, Inc. 8310 16<sup>th</sup> Street, MS 486 Sturtevant, WI 53177-0902</b> <input type="checkbox"/> Check if this is a new address	6. <b>Expedited Review.</b> In accordance with FIFRA Section 3 (c). (3) (b) (i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional Page(s) if necessary. (For section I and Section II)

Notification to add NSF logo to label.

Contact information: kresti.Lyddon@johnsondiversey.com or Kresti A. Lyddon, fax: 262-631-4036

### Section - III

#### 1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt. _____ No. per Container _____	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Package wgt. _____ No. Per Container _____	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
--	--	---	---

\* Certification must be submitted

3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container <b>1, 5.12 55 &amp; 330 gallons bulk</b>	5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application.)		
Name <b>Kresti A. Lyddon</b>	Title <b>Senior Registration Specialist</b>	Telephone No. (Include Area Code) <b>262.631.4830</b>

#### Certification

I certify that the statements which I have made on this form and all attachments are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received (Stampcc)

2. Signature 	3. Title <b>Senior Registration Specialist</b>	6. Date Application Received (Stampcc)
4. Typed Name <b>Kresti A. Lyddon</b>	5. Date <b>October 24, 2007</b>	



70  
JohnsonDiversey, Inc.  
8310 16th Street  
P.O. Box 902  
Sturtevant, WI 53177-0902

4/1/0  
JohnsonDiversey  
Clean is just the beginning



October 24, 2007

Document Processing Desk (NOTIF)  
Attn: Emily Mitchell, Antimicrobial Division  
Office of Pesticide Programs (7510C)  
U. S. Environmental Protection Agency  
Room 266A, Crystal Mall 2  
1801 South Bell Street  
Arlington, VA 22202

COURIER

Dear Ms. Mitchell:

Subject: Label Notification for Dibac™  
EPA Reg. No. - 875-47

JohnsonDiversey, Inc. is submitting some minor changes to the label for Dibac™, EPA Reg. No. 875-47. We are proposing to add the NSF logo, as our application for NSF listing of this product with the National Sanitation Foundation has received their approval. Enclosed are the following documents in support of this notification:

- Application for Pesticide Registration (EPA Form 8570-1)
- Three copies of proposed label
- Certification of Notification per PR Notice 98-10


If you have any questions regarding this application or require further information, please contact me at 262-631-4830 or at [kresti.lyddon@johnsondiversey.com](mailto:kresti.lyddon@johnsondiversey.com)

Sincerely,



Kresti A. Lyddon  
Senior Registration Specialist

Enclosures



5/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

JohnsonDiversey



DIBAC™

Liquid Chlorinated Sanitizer

Liquid Bactericide • Disinfectant • Sanitizer • Deodorizer

For Industry (Institutional) Use (Only)

<b>ACTIVE INGREDIENT:</b>	
Sodium Hypochlorite.....	9.2%
<b>OTHER INGREDIENTS:</b> .....	90.8%
<b>TOTAL:</b> .....	100.0%

Available Chlorine: 8.74%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE.**

**1-800-851-7145**

*Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.*

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

See additional precautionary statements on back (side) (left) (right) (panel) (of) (label) (below).

(See reference sheet (enclosed in case) for additional directions for use.)

Net Contents:

(Product of USA)

6/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Note: This product degrades with age. Use a chlorine test kit and increase/decrease dosage, as necessary to obtain the required level of available chlorine.

**Concentration Chart**

<u>Available Chlorine</u>	<u>Dilution Rate</u>
25 ppm	1 oz. to 20 gallons
50 ppm	1 oz. to 10 gallons
100 ppm	2 oz. to 10 gallons
200 ppm	4 oz. to 10 gallons
500 ppm	10 oz. to 10 gallons

**CLEAN-IN-PLACE SANITIZING OF EQUIPMENT, PIPELINES AND FILLERS:**

Thoroughly flush, clean and potable water rinse system. Prepare a volume equal to 110% of system capacity of 200 ppm available chlorine sanitizer by mixing 4 oz. of product to 10 gallons of water. Pump sanitizer through the system until full flow is obtained at all extremities and the system is completely filled with the sanitizer and all air is removed. Close drain valves and hold under pressure for at least 10 minutes to insure contact with all internal surfaces. Remove some sanitizer solution from a drain valve and test with a Chlorine Test Kit. If the sanitizer contains less than 50 ppm available chlorine, repeat the entire cleaning/sanitizing process. Completely drain and air-blow the system to remove residual sanitizer before restarting processing.

**RESTAURANTS, INSTITUTIONS, AND OTHER FOOD SERVING ESTABLISHMENTS:**

1. Scrape and pre-wash utensils, dishes and glasses to remove gross soil.
2. Wash with recommended detergent.
3. Rinse with clean (potable) water.
4. Sanitize in a solution containing 4 oz. of product for each 10 gallons of tap water used (200 ppm available chlorine). Immerse all utensils for at least 2 minutes or contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.

Note: NO POTABLE WATER RINSING of equipment is required. Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitization purposes. Test sanitizer frequently during operation with a Chlorine Test Kit to insure that solution does not drop below 50 ppm available chlorine. If no test kit is available or if available chlorine has dropped below 50 ppm during sanitizing, either discard the sanitizer solution or add sufficient of product to reestablish a 200 ppm available chlorine sanitizer strength.

**SANITIZING OF POROUS FOOD CONTACT SURFACES:**

**Meat, Beverage and Food Processing Plants** - Clean and potable water rinse cutting boards, plastic baskets and rubber gaskets. Sanitize by preparing a solution of 600 ppm available chlorine sanitizer by mixing 12 oz. of product in 10 gallons of water. Rinse, flood, immerse, wipe or spray porous surfaces until thoroughly wet. Maintain contact with the sanitizer for at least 2 minutes. Drain excess sanitizer. Prior to reusing treated equipment, rinse in a fresh sanitizer solution containing 200 ppm available chlorine by mixing 4 oz. of product in 10 gallons of water. Drain excess sanitizer. Do not rinse and do not soak equipment overnight. Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitization purposes.

7/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

**SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES:**

**Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSE OR FLOOD APPLICATIONS.** Prepare a sanitizing solution by thoroughly mixing 6 oz of this product with 5 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse, immerse or flood all surfaces thoroughly with the sanitizing solution, maintaining contact with the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

**SANITIZATION OF NON-POROUS FOOD CONTACT SURFACES:**

**Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSE OR FLOOD APPLICATIONS.** A solution of 100 ppm available chlorine may be used in the sanitizing solution if a test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2 oz of this product in 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 4 oz of this product in 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If the solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitization purposes.

**SANITIZATION OF NON-POROUS NON-FOOD CONTACT SURFACES:**

**Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSE OR FLOOD APPLICATIONS.** Prepare a sanitizing solution by thoroughly mixing 2 oz of this product with 5 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the sanitizing solution, maintaining contact with the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

**DISINFECTION OF NON-POROUS NON-FOOD CONTACT SURFACES:**

**Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSE OR FLOOD APPLICATIONS.** Prepare a disinfecting solution by thoroughly mixing 12 oz of this product with 10 gallons of water to provide approximately 600 ppm available chlorine. Clean equipment in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the disinfecting solution, maintaining contact with the solution for at least 10 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

**SPRAY AND FOGGING SANITIZATION METHOD:**

Pre-clean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the production in a ratio of 4 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 12 oz. per 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with a 600 ppm solution with a 200 ppm solution.

8/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

**MECHANICAL WASHERS:**

**Wash-Dump or Sanitizing Rinse Section Types** - After cleaning and rinsing equipment or containers, apply a sanitizing spray rinse containing 100 ppm available chlorine by adjusting mechanical feeding device to meter 2 oz. of product per 10 gallons of water. Test sanitizer frequently during operation with a Chlorine Test Kit to insure that solution does not drop below 50 ppm available chlorine. Remove and drain sanitized items. Sanitizer used in automatic systems may be used in the general cleaning process but may NOT be re-used for sanitizing purposes.

If no test kit is available or if available chlorine has dropped below 50 ppm during sanitizing, either discard the sanitizer solution or add sufficient quantity of product to reestablish a 200 ppm available chlorine sanitizer strength. Resanitize equipment. Do not rinse equipment with water after sanitizing and do not soak equipment overnight in sanitizer solution.

**POST-HARVEST PROTECTION** – Potatoes can be sanitized after cleaning and prior to storage by spraying with a sanitizing solution at a level of 1 gallon of sanitizing solution per ton of potatoes. Thoroughly mix 2 oz of this product to 2 gallons of water to obtain 500 ppm available chlorine.

**FRUIT AND VEGETABLE WASHING** – Thoroughly clean all fruit and vegetables in a tank or spray washer. Dilute 1 oz. of product in 20 gallons of water to make a sanitizing solution containing 25 ppm available chlorine. Drain and rinse product with potable water. Immerse or spray in a separate wash tank for 2 minutes. Check available chlorine content with test kit. Spray rinse vegetables with additional sanitizer only prior to packaging. Rinse fruit and vegetables with potable water.

**FOOD EGG SANITIZING** – Thoroughly clean all eggs. Dilute 4 oz. of product for each 10 gallons of warm water to produce 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130 F. Spray the sanitizer so that all eggs are thoroughly wetted. Allow the eggs to completely dry before casing or breaking. Do not apply a potable water rinse. Do not reuse solution for sanitizing eggs.

**DISINFECTION OF NON-POROUS NON-FOOD CONTACT SURFACES:**

**Meat, Poultry and Food Processing Plants** - Clean and rinse pallets, bins, walls, floors, refrigeration areas and other non-food contact surfaces. Prepare a disinfecting solution by mixing 12 oz. of product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Prior to use, expose equipment and surfaces to disinfecting solution by rinsing, immersion, wiping or spraying to thoroughly wet surface. Allow solution to remain in contact for at least 10 minutes. Drain excess disinfectant from surfaces. Do NOT rinse equipment with water after treatment and do not soak equipment overnight.

**DISINFECTION OF DRINKING WATER:**

Prepare a stock disinfectant solution by mixing 1 oz. of product with 5 gallons water to obtain 100 ppm available chlorine. Meter the stock solutions with hypochlorinator until a free residual chlorine content of at least 0.2 ppm but no more than 0.6 ppm is obtained throughout the distribution system. Check water frequently with a Chlorine Test Kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Primary Drinking Water Regulation. Contact your local Health Department for further details.

**SANITIZING POROUS FOOD CONTACT EQUIPMENT:**

Prepare a 600 ppm solution by thoroughly mixing 12 ounces of this product in 10 gals water. Clean surfaces in the normal manner. Rinse, immerse or flood all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution (4 oz / 10 gallons). Do not rinse and do not soak overnight.



9/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

**SANITIZATION OF CONVEYORS FOR MEAT, POULTRY, SEAFOOD, FRUITS, AND VEGETABLES**

For use in the static or continuous washing, rinsing and sanitizing of conveyor equipment, peelers, collators, slicers, saws, etc.

1. Remove all products from equipment if during treatment the sanitizer will directly contact the items.
2. Prepare 200 ppm solution of this product (4 oz / 10 gallons).
3. Apply sanitizer solution to the return portion of the conveyor or to the equipment using a coarse spray, other means of wetting the surfaces. Treat for at least one (1) minute. Control the volume of solution so as to permit maximum drainage and to prevent puddles. The conveyor may still be damp when food contact occurs.
4. Allow equipment to drain adequately before reusing; a dry surface is not required.

**TREATMENT OF POULTRY PROCESSING WATER**

Follow guidelines of local water authority for water potability treatment.

**Continuous Feed:** Using an automatic metering device, continuously feed this product into the water to obtain and/or maintain a level of 5-20 ppm available chlorine (1.2 oz product per 100 gal. water, to 4 oz product per 100 gals. Water). Confirm target chlorine level with either a chlorine test kit or an automatic testing device. When the available chlorine level reaches 10 ppm, notify the USDA plant inspector.

**Intermittent Feed:** Start up by adding 2 oz of product per 1000 gallons of water for each 1 ppm of available chlorine needed. For subsequent doses, check chlorine level with a chlorine test kit. Add enough of this product to maintain the target chlorine level and confirm this level with a chlorine test kit. Do not pour this product directly on poultry product in the water.

**COOLING TOWER/EVAPORATIVE CONDENSER WATER: CONTINUOUS FEED METHOD:**

**Initial Dose:** When system is noticeably fouled, apply 58 to 116 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. **Subsequent Dose:** Maintain this treatment level by starting a continuous feed of 1 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

**COMMERCIAL LAUNDRY SANITIZERS:** Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 4 oz. of this product with 10 gallons of water to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine if solution has been allowed to stand. Add more of the product if the available chlorine level has dropped below 200 ppm.

**FARM PREMISES:** Remove all animals, poultry, and feed from premises, vehicles and enclosures. Remove all litter, and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or transverse by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 20 oz of this product with 10 gallons of water. Immerse all halters, ropes and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, cars, boats and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be rinsed with potable water before use.

**HAND SANITIZING:** Thoroughly wash hands with soap or detergent. Rinse with potable water. Prepare a solution containing 1 oz of this product per 10 gallons of water providing 50 ppm available chlorine (use test kit to determine exact available chlorine). Submerge hands. Let air dry.

**SHOE BATH SANITIZER DIRECTIONS:** To prevent tracking harmful organisms into animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made use solution should be placed at all entrances to buildings, hatcheries and all entrances to the production and packaging rooms. Scrape waterproof shoes and place them into a solution containing 0.25 ounces of this product per gallon of water (or equivalent use dilution)(200 ppm active solution) for 60 seconds prior to entering area. Change use-solution in the bath daily or sooner if use-solution appears dirty.

10/10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### ENVIRONMENTAL HAZARDS (for containers of 5 gallons or more)

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

## PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**DANGER: Corrosive.** Causes irreversible eye damage and skin burns. Harmful if swallowed. Wear chemical splash-proof goggles or face shield, rubber gloves and protective clothing when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

### PHYSICAL OR CHEMICAL HAZARDS

**STRONG OXIDIZING AGENT:** Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release irritating chlorine gases which are harmful to eyes, lungs and mucous membranes.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

### PELIGRO

Si no puede leer en ingles, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto.

EPA Reg. No. 875-47

EPA Est. No.

Lot code indicates establishment number.

© 2007 JohnsonDiversey, Inc., Sturtevant, WI 53177 U.S.A. All Rights Reserved.

