

WEEDS CONTROLLED

BANVEL Herbicide, when applied in accordance with this label, will give control of many ANNUAL and PERENNIAL broadleaf weeds including:

ANNUALS

amaranth, spiny	henbit	purslane, horse
beggarweed, Florida	jimsonweed	pusley, Florida
burclover, California	lambsquarters, common	ragweed, common
buttercup (<i>Ranunculus</i> spp.)	lettuce, prickly	ragweed, giant
chickweed, common	marshcress	rocket, London
clovers, annual	morningglory, ivyleaf	shepherdspurse
cocklebur, common	morningglory, tall	sicklepod
cudweed, wandering	mustards, wild	sida, prickly
(rabbit tobacco)	nightshade, black	(teaweed)
eveningprimrose, cutleaf	pepperweed, Virginia	speedwell, corn
fireweed	pigweed, prostrate	speedwell, purslane
(American burnweed)	pigweed, redroot	sowthistle, annual
fleabane, annual	(carelessweed)	spurge, prostrate
geranium, Carolina	pigweed, smooth	swinecress
groundcherry, smooth	purslane, common	toadflax, oldfield
		woodsorrel, southern yellow

PERENNIALS

*aster spp.
 chickweed, mouseear
 dandelion, common
 dock, curly
 dogfennel
 elderberry
 (American elder)
 fleabane, Philadelphia
 garlic, wild
 *goldenrod spp.
 groundsel, cressleaf
 onion, wild
 sowthistle, perennial

* Noted PERENNIAL weeds require higher rates than other listed PERENNIAL species. See APPLICATION RATES AND TIMINGS section of this label on pages - .

APPLICATION RATES AND TIMINGS

Application of BANVEL Herbicide can be made any time after the weeds have emerged but prior to the close-in stage of sugarcane, Application rates and timings are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

- For control of listed ANNUAL broadleaf weeds, apply 1/2 to 1 pint (1/4 - 1/2 lb. a. i.) per treated acre of BANVEL Herbicide when weeds are less than 4 inches and actively growing. Apply 1 to 1 1/2 pints (1/2 - 3/4 lbs. a. i.) per treated acre when treating established weed growth.

- 7
000
- For control of listed PERENNIAL broadleaf weeds, apply 1 1/2 to 3 pints (3/4 - 1 1/2 lbs. a.i.) per treated acre of BANVEL Herbicide to the foliage of actively growing weeds.
 - For control of noted * PERENNIAL broadleaf weeds, apply 3 to 6 pints (1 1/2 - 3 lbs. a.i.) per treated acre.

Retreatments may be made as needed, however, do not exceed 3 quarts (3 lbs. a.i.) per treated acre of BANVEL Herbicide during a growing season.

TANK MIX TREATMENT

BANVEL Herbicide may be tank mixed with 2,4-D for control of undesirable vegetations listed in this label (refer to pages and) as well as additional weeds listed on the 2,4-D product label.

READ AND FOLLOW 2,4-D PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS AND RESTRICTIONS.

- For control of ANNUAL broadleaf weeds, tank mix 1/2 to 1 pint (1/4 - 1/2 lb. a.i.) of BANVEL Herbicide with 1/2 to 3/4 lb. acid equivalent of 2,4-D per treated acre.
- For control of PERENNIAL broadleaf weeds, tank mix 1 to 1 1/2 pints (1/2 - 3/4 lb. a.i.) of BANVEL Herbicide with 3/4 to 1 1/2 lbs. acid equivalent of 2,4-D per treated acre. This treatment may be used for more effective control of established ANNUAL broadleaf weeds.
- For control of difficult to control weeds such as the noted * PERENNIAL broadleaf weeds, tank mix 1 1/2 to 3 pints (3/4 - 1 1/2 lbs. a.i.) of BANVEL Herbicide with 1 1/2 to 3 lbs. acid equivalent of 2,4-D per treated acre.