ACCEPTED

AUG 1 1 ...

... REGISTERED UNDER SEC. 3(c)(5) FOR THE PRODUCT NO. 876-37 SUBJECT TO THE ATTACHED COMMENTS.

# VELSICOL®

# BANVEL-D 4-S

## HERBICIDE

**ACTIVE INGREDIENTS:**

| | |
|---|---|
| Dimethylamine salt of dicamba (3, 6-dichloro-o-anisic acid)* | 49.0% |
| Dimethylamine Salts of related acids | 7.9% |
| **INERT INGREDIENTS** | **43.1%** |
| TOTAL | 100.0% |

*Equivalent to 40.6% 3, 6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon.

Contains 4 pounds dicamba equivalent per gallon.

## CAUTION:

**Keep out of reach of children.**

**(See left panel for additional caution statements)**

**CONTENTS: 5 GALLONS**

**VELSICOL CHEMICAL CORPORATION • 341 East Ohio • Chicago, Illinois 60611**

**CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.**

**IMPORTANT: Avoid application on or near desirable trees, or other plants, or on areas where their roots may extend or in location where the chemical may be washed or moved into contact with their roots. Do not clean equipment on or near desirable trees or other plants. Prevent drift of product to desirable plants. Avoid contamination of domestic or irrigation waters. Thoroughly clean application equipment after use.**

**USDA Registration Number 876-37**

**DIRECTIONS: For use on lawns, turf (not pastures) and fairways, aprons, tees and rough.**

| WEEDS | | DOSAGE | APPLICATION |
|---|---|---|---|
| curly dock<br>common chickweed<br>mouseear chickweed<br>stitchwort<br>dog fennel<br>knawel<br>chicory<br>henbit | English Daisy<br>Spurge<br>Purslane<br>Pepperweed<br>Carpetweed<br>Spurry | 1-2 pints Banvel D-4S per acre or 2-4 teaspoons per 900 square feet (½ to 1 lb. Dicamba per acre) | Apply as a foliar spray to actively growing weeds with sufficient water to give good coverage. Fall germinating weeds may be controlled more effectively when treated in the fall or early winter. Make only one application per year. Banvel D may be used on newly seeded turf after second mowing. |
| **DIRECTIONS: For use on BENTGRASS GREENS.** | | **DOSAGE** | |
| curly dock<br>common chickweed<br>mouseear chickweed<br>stitchwort | dog fennel<br>knotweed<br>sheep sorrel<br>clover | ½ to 1 pint Banvel D-4S per acre or 1-2 teaspoons per 900 square feet. (¼ to ½ pound Dicamba per acre) | |

DIRECTIONS: For use on roadways (not turf or pastures).

| WEEDS | DOSAGE/ACRE | APPLICATION |
|---|---|---|
| Field bindweed, Russian knapweed, Canada thistle, Leafy spurge | 1 to 2 gallons (4 to 8 lbs. dicamba acid equivalent) | Apply to foliage and/or soil when weed is in active growing stage. For spot treatment mix 1-1/2 to 3 tablespoons with one gallon of water to treat one square rod (272 sq. ft.) |
| Top growth control of field bindweed, Russian knapweed, Canada thistle, and Leafy spurge. For control of chickweeds (common, mouseear, perennial or stitchwort), curly dock, dog fennels, (mayweed and corn chamomile), knapweed, clover, sheep sorrel, (red sorrel) | 1 to 2 pints ( 8 to 16 oz. dicamba acid equivalent) | Apply as a foliar spray to actively growing weeds. |

**NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.**

**Container Disposal: Do not reuse empty container. Crush or perforate and bury empty container.**