

VELSICOL®

Banvel® 4-S

HERBICIDE

ACCEPTED
JUL 1980
Under the
Fungicide
as amended
registration
EPA Reg. No. 876-37

E.P.A. REG. NO. 876-37-AA
E.P.A. EST. NO. 876-TX-2

Net Contents:
2½ Gallons

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3, 6-dichloro- <i>o</i> -anisic acid)	48.2%
Dimethylamine Salts of related acids ..	12.0%
INERT INGREDIENTS	39.8%
TOTAL	100.0%

Equivalent to 40.0% 3,6-dichloro-*o*-anisic acid (dicamba) or 4 pounds per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional caution statements

VELSICOL CHEMICAL CORPORATION
341 E. 310 ST., CHICAGO, ILL. 60611

79

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid application on or near desirable trees, or other plants or on areas where their roots may extend or in location where the chemical may be washed or moved into contact with their roots. Do not clean equipment on or near desirable trees or other plants. Prevent drift of product to desirable plants. Coarse sprays are less likely to drift. Avoid contamination of domestic or irrigation waters. Thoroughly clean application equipment after use.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

CONTAINER DISPOSAL: Do not reuse empty container. Crush or perforate and bury empty container.

LIMIT OF WARRANTY AND LIABILITY

The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances and from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

07152-205AB

DIRECTIONS: For use on lawns, turf (not pastures) and fairways, aprons, tees and rough.

WEEDS	DOSAGE	APPLICATION
curly dock common chickweed mouseear chickweed stichwort dog fennel knapweed sheep sorrel clover knapweed chicory henbit english daisy spurge purslane pepperweed carpetweed lawn burweed hawkweed spurry	1-2 pints Banvel 4-S per acre or 2-3 teaspoons per 800 square feet (1/2 to 1 lb. Dicamba per acre.)	Apply as a foliar spray to actively growing weeds with sufficient water to give good coverage. Fall germinating weeds may be controlled more effectively when treated in the fall or early winter. Make only one application per year. Banvel may be used on newly seeded turf after second mowing.

DIRECTIONS: For use on BENTGRASS GREENS.

curly dock common chickweed mouseear chickweed stichwort dog fennel knapweed sheep sorrel clover	1/2 to 1 pint Banvel 4-S per acre or 1-2 teaspoons per 800 square feet. (1/2 to 1/2 pound Dicamba per acre)	
---	---	--

DIRECTIONS: For use on roadways (not turf or pastures).

WEEDS	DOSAGE/ACRE	APPLICATION
Field bindweed, Russian knapweed, Canada thistle, Leaky spurge.	1 to 2 gallons (4 to 8 lbs. Dicamba acid equivalent).	Apply to foliage and/or soil when weed is in active growing stage. For spot treatment mix 1 1/2 to 3 tablespoons with one gallon of water to treat one square rod (272 sq. ft.).
Top growth control of field bindweed, Russian knapweed, Canada thistle, and Leaky spurge. For control of chickweeds (common, mouseear, perennial or stichwort), curly dock, dog fennel (mayweed and corn chamomile), knapweed, clover, sheep sorrel, (red sorrel).	1 to 2 pints (8 to 16 oz. Dicamba acid equivalent).	Apply as a foliar spray to actively growing weeds.

90