

**WARNING:** May Produce Delayed Chemical Burns. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Keep children and pets off treated areas until dry. Use only according to directions.

**This product is toxic to fish and wildlife.**

**Birds and other wildlife in treated areas may be harmed.**

**ANTIDOTE:** If swallowed. Give milk or white of egg beaten with water, then a tablespoonful of salt in a glass of warm water, and repeat until vomit fluid is clear. Repeat milk or white of egg beaten with water. Call a Physician immediately.

**If on skin.** Wash immediately with soap and warm water. **If in eyes.** Flush with water for at least 15 minutes, and get medical attention.

**CONTAINER DISPOSAL:** Decontaminate by repeated rinsing with water. Crush or perforate and bury when empty. NEVER REUSE.

DIRECTIONS: Turf and Lawn Diseases:			
Diseases	grasses	DOSAGE/1,000 square feet	application
Dollar Spot Brown Patch Copper Spot Melting Out	Bermuda grass St. Augustine Centipede Kentucky Bluegrass Merion Bluegrass Bentgrass	one ounce of Velsicol Memmi .8EC to 3-5 gallons of water per 1,000 square feet. Use twice the amount of water at temperatures over 85°F.	Repeat application at 10 to 14 day intervals if necessary. Make no more than 2 applications in a 30 day period to Bentgrass areas.
gray leaf spot	St. Augustine grass	1.5 ounces of Velsicol Memmi .8EC to 3-5 gallons of water to treat 1,000 square feet.	Repeat application at 10 to 14 day intervals if necessary.
cottony blight (Pythium)	ryegrass	two ounces of Velsicol Memmi .8EC to 3-5 gallons of water to treat 1,000 square feet. Use twice the amount of water at temperatures over 85°F.	Repeat application at 7 to 10 day intervals if necessary. Do not less than 30 days before or after overseeding with ryegrass.

Gladiolus Corms

Diseases:	application
Scab Fusarium Curvularia	mix 1 pint of Velsicol MEMMI .8EC in 50 gallons of water. Treat two batches (volume of corms that can be thoroughly covered by the solution) of corms and allow the solution to drain from the corms back into the treating tank each time, and then renew the solution by adding ½ pint of Velsicol MEMMI .8EC. Treat two more batches of corms and discard the solution. Soak corms two hours. Plant corms the same day as treated.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty of any kind, either express or implied, and does not warrant results nor residues no greater than any tolerance which may be established by appropriate governmental agencies, nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, misuse, handling or storage of this material, whether in accordance with directions or not.

FORM 400-130 R4 PRINTED IN U.S.A. CG 4/67

ACCEPTED

OCT. 31, 1967

U.S. DEPT. OF AGRICULTURE FEDERAL INSECTICIDE  
FERTILIZER AND SOIL CONDITIONER ACT  
POISON REGISTERED  
876-38

**MERCURY FUN**

Active Ingredients:  
N-Methylmercuri-1  
tetrahydro-3, 6-e  
ano-3, 4, 5, 6, 7  
chlorophthalimide  
(Mercury Content  
Inert Ingredients . . .  
Total . . . . .

ACCEPTED WITH COMMENTS

UNION FEDERAL INSECTICIDE  
FUNGICIDE AND FUNGICIDE ACT  
FOR ECONOMIC PESTICIDE PROTECTION

Oct. 31, 1947

ACCEPTED

# VELSICOL®

## MEMMI .8EC

**DANGER**

**POISON**  **POISON**

### MERCURY FUNGICIDE

Active Ingredients:

N-Methylmercuri-1, 2, 3, 6-tetrahydro-3, 6-endomethano-3, 4, 5, 6, 7, 7-hexachlorophthalimide . . . 10.0%  
(Mercury Content 3.4%)

Inert Ingredients . . . . . 90.0%

Total . . . . . 100.0%

**KEEP OUT OF THE REACH OF CHILDREN**

See Left Panel for Antidote and Additional Precautionary Statements.

Contents: 1 Gallon



**VELSICOL® CHEMICAL CORPORATION 341 East Ohio Street, Chicago, Illinois 60611**

U.S.D.A. Reg. No. 876-38