

# VELSICOL<sup>®</sup>

## Banvel<sup>®</sup> D+2,4-D

(Contains 1.5 Pounds Dicamba and 3 Pounds 2,4-D)

CONTENTS: 5 Gallons

### CAUTION

**Keep out of the reach of children.  
See left panel for additional cautions.**

**VELSICOL CHEMICAL CORPORATION**  
341 East Ohio St., Chicago, Illinois 60611



ACCEPTED

OCT. 19, 1967


UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROBERTSON ACT  
FOR THE AGRICULTURAL DIVISION REGISTERED  
ED UNDER NO. 876-42 SUBJECT  
TO ATTACHED COMMENTS.

#### Active Ingredients:

Dimethylamine Salt of dicamba (3, 6 dichloro-o-anisic acid)*	18.4%
Dimethylamine Salts of related acids	3.2%
Dimethylamine Salt of 2,4-dichloro- phenoxyacetic acid**	36.8%
Inert Ingredients	41.6%
Total	100.0%

\*Equivalent to 15.2% 3,6 dichloro-o-anisic acid (dicamba) or 1.5 pounds per gallon.

\*\*Equivalent to 30.4% 2,4 dichlorophenoxyacetic acid or 3 pounds per gallon.



**CAUTION:** Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

**IMPORTANT:** Avoid drift of spray mist onto vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not apply within the drip line of trees or shrubs. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizers, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

USDA Registration Number 876-42

USE	WEEDS	DOSAGE	APPLICATION
On lawns, turf (not pastures), and golf fairways, aprons, tees and rough.	dandelion <del>plantain</del> curly dock common chickweed mouseear chickweed stitchwort dog yennel sheep sorrel clover knawel chicory	henbit ragweed <del>pigweed</del> knotweed ground ivy (creeping charlie) wild onion burdock wild garlic poison oak poison ivy	Apply as a foliar spray to actively growing weeds with sufficient water to give good coverage. Fall germinating weeds may be controlled more effectively when treated in the fall or early winter. Make only one application per year.
	English daisy <del>spurge</del> purslane pepperweed	lawn burweed hawkweed spurry cat's paw	
		Apply 2 pints to 3 pints BANVEL D + 2,4-D per acre or 4 to 6 teaspoons per 1000 square feet. (Equivalent to .34 lb. Dicamba + .68 lb. 2,4-D to .51 Dicamba + 1.02 lbs. 2,4-D per acre).	
		Apply 3 pints BANVEL D + 2,4-D per acre or 6 teaspoons per 1000 square ft. (Equivalent to .51 lb. Dicamba + 1.02 lbs. 2,4-D per acre).	

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty of any kind, either express or implied, and does not warrant results nor residues no greater than any tolerance which may be established by appropriate governmental agencies, nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, misuse, handling or storage of this material, whether in accordance with directions or not.

FORM 400-144R3 PRINTED IN U.S.A. CG 966

ACCEPTED  
Oct. 17, 1967

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROBBERS ACT  
FOR ECONOMIC PEST CONTROL  
E.P. No. 876-42  
TO APPLICANTS