

VELSICOL®

Banvel® K

HERBICIDE

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See Back Panel for Additional Cautions

**E.P.A. REG. NO. 876-44-AA
E.P.A. EST. NO. 876-TX-2**

**Net Contents:
2½ Gallons**

ACTIVE INGREDIENTS:

Dimethylamine Salt of dicamba
(3,6-dichloro-o-anisic acid)* 15.5%
Dimethylamine salt of
related acids 3.9%
Dimethylamine salt of 2,4-dichlorophenoxy
acetic acid** 24.7%

INERT INGREDIENTS: 50.9%
TOTAL 100.0%

*Equivalent to 12.9% 3,6-dichloro-o-anisic acid.

**Equivalent to 25.9% 2,4-dichlorophenoxyacetic acid.

Contains: 1.25 Lbs. Dicamba and 2.34 Lbs. 2,4-D
Per Gallon

**VELSICOL CHEMICAL CORPORATION
341 E. OHIO ST., CHICAGO, ILL. 60611**

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinsate that cannot be used according to label directions must be disposed of according to Federal, State, or local procedures under the Resource Conservation and Recovery Act

3. CONTAINER DISPOSAL

Plastic

Triple rinse (or equivalent) and dispose of in a sanitary landfill or by incineration if permitted by State and local authorities

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions, or from mixing this product before use with any substance except as recommended by the product's label; (ii) arising from handling or storage in violation of label instructions; (iii) for all indirect special or consequential damages; (iv) when not reported to this company within one year of discovery; and (v) arising from product not used within the label-designated shelf life or four years from date of purchase, whichever first occurs. THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

Practical Treatment:

External:

For Skin: Wash with mild soap and water. Rinse with fresh water.

For Eyes: Rinse eyes with water for at least 15 minutes. Get medical aid.

Internal:

If swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with a finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled: Remove victim to fresh air. Apply respiration if necessary.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on label.

IMPORTANT

Do not apply BANVEL® K Herbicide directly to or allow spray drifts to come in contact with desirable plants, particularly beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf crops. Coarse sprays are less likely to drift.

Do not contaminate domestic or irrigation water by cleaning of equipment or disposal of wastes.

Do not store in direct contact with fertilizer, seed, insecticides, or fungicides.

Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes.

Do not graze or harvest for dairy or beef feed prior to ensilage stage (milk stage).

Do not apply (except as recommended) or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemicals may be washed or moved into contact with their roots.

The use directions for this product are intended for crops grown under normal growing conditions. Crops grown under stress such as drought, insect damage, or frost may develop injury from the use of this product.

Consult your State Agricultural Experiment Station or Extension Service weed specialists for advice concerning special local use situations or tolerance of specific crop varieties to BANVEL K Herbicide.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read LIMIT OF WARRANTY AND LIABILITY before buying or using. If terms are not acceptable, return at once unopened.

BANVEL® K SPRAY DRIFT: Soybeans, tobacco and other broadleaf crops are sensitive to BANVEL K as it requires only a small amount of chemical per acre for symptoms to appear. Symptoms generally appear about two weeks after drift occurs. Soybeans generally outgrow these initial symptoms without any adverse effect on soybean yields. Affected tobacco leaves will be permanently malformed.

To minimize BANVEL K spray drift onto soybeans and other sensitive crops, observe the following precautions:

- 1 Use coarse sprays. Use 20 gallons or more of water per acre. Keep the spray pressure below 20 psi. For broadcast application, use TK5 tips (or equivalent).

- 2 Spray with the nozzles as close to the weeds as is practical for good coverage. When susceptible crops are nearby, do not apply BANVEL K Herbicide after corn is 24 inches tall.

- 3 Do not apply BANVEL K Herbicide to field corn when the daily temperature is expected to be over 85° F.

- 4 Make application when there is some air movement but wind is less than 5 miles per hour. If any wind over 5 mph is blowing in the direction of sensitive crops, do not spray.

Do not apply BANVEL K Herbicide in the vicinity of soybeans if any one of the following conditions exists:

- a The corn is more than 24 inches tall.
- b Soybeans in the vicinity are more than 10 inches tall.
- c Soybeans in the vicinity have begun to bloom.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Fall seeded wheat (not underseeded to legumes)

WEED: For the control of Hard to Kill weeds (phenoxy resistant) including wild buckwheat, ladythumb, smartweed, knawel, corn cockle, mayweed, corn chamomile.

And for the control of other broadleaf weeds including Canada thistle (top growth), perennial sowthistle (top growth), lambsquarters, pigweed, cocklebur, dragonhead (mint), mustards (annual), kochia, hempnettle, wild radish, wild garlic (top growth), velch, Russian thistle, ragweed, stinkweed, goatsbeard (salsify), dandelion, Frenchweed (field pennycress), puncturevine, field peppergrass (field pepperweed), sunflower and others.

APPLICATION RATE BANVEL® K HERBICIDE: Apply 0.8 pint BANVEL® K Herbicide (1/3 lb dicamba plus 1/3 lb 2,4-D) per acre.

One gallon treats 10 acres.

APPLICATION DIRECTIONS: Apply as a foliar spray immediately after winter dormancy and before the grain begins to joint.

Apply when weeds are actively growing.

Apply in 3 1/2 to 40 gallons of water per acre.

Field corn (postemergence)

WEED: For the control of Hard to Kill weeds (phenoxy resistant) including field bindweed, Canada thistle, perennial sowthistle, horsenettle, prostrate spurge, smartweed, common milkweed, hedge bindweed.

And for the control of other broadleaf weeds including buckthorn, burdock, buttonweed, clover, cocklebur, flower-of-an-hour, jimsonweed, lambsquarters, common morningglory, mustard, pepperweed, pigweed, poorjoe, ragweed, spanishneedles, velvetleaf, waterhemp.

APPLICATION RATE BANVEL® K HERBICIDE: Broadcast (overall or directed) 0.8 to 1.6 pint (1/3 to 1/2 lb dicamba + 1/3 to 1/2 lb 2,4-D) per acre.

One gallon treats 5-10 acres.

Band (overall or directed) 1/3 of total area) 0.26 to 0.52 pint (2/3-1 oz dicamba + 1/3-2/3 oz 2,4-D) per acre. One gallon treats 15-30 acres.

APPLICATION DIRECTIONS: May be applied as a post-emergence treatment over the tops of the corn plant (overall) on corn less than 8 in height.

When corn is 8 or more in height, use drop nozzles and direct spray at base of corn plant.

Do not apply after corn is 36 inches high.

Make only one application per season.

DO NOT GRAZE TREATED FIELDS OR HARVEST FOR DAIRY OR BEEF FEED PRIOR TO ENSILAGE STAGE (milk stage).