

ak®

CAUTION

Avoid contact with skin, eyes, or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention.

IMPORTANT

Do not apply MONDAK® herbicide directly to or allow spray drifts to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, sugar beets, beans, or other desirable plants. (Coarse sprays are less likely to drift.) Do not contaminate domestic or irrigation water. Do not store in direct contact with fertilizer, seed, insecticides or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. When container is empty, immediately wash thoroughly and destroy. Do not graze treated areas or harvest for dairy feed prior to crop maturity. Do not apply (except as recommended) or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

LIMIT OF WARRANTY AND LIABILITY

The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances and from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

DIRECTIONS FOR USE

WHEAT (not underseeded to legumes)

For the control of "hard to kill" weeds (phenox-resistant) including:

Bedstraw	Henbit
Corn cockle	Knawel (German moss)
Corn cockle	Ladysthumb
Corn buttercup (crowfoot)	Miner's lettuce
Big larnet	Prickly lettuce
Field pennycress (farweed)	Smartweed
Groundsel	Wild Buckwheat

And for the control of other broadleaf weeds including:

Canada thistle (top growth)	Perennial Sowthistle (top growth)
Cocklebur	Pigweed (redroot)
Dandelion	Puncturevine
Dragonhead (mint)	Ragweed (common)
Field pennycress (farweed)	Russian thistle
Field pepperweed	Shepherdspurse
Hempnettle	Sunflower
Knotweed	Vetch
Kochia	Wild radish
Lambsquarters	and others
Mustard (annual)	

For ground application use 5 to 20 gallons of water per acre.

For aerial application use 3 to 8 gallons of water per acre.

Spring Seeded Wheat: For the control of the listed weeds, apply 0.8 pint of MONDAK® herbicide (1/2 pound of dicamba plus 1/4 pound of MCPA) per acre. One gallon of MONDAK® herbicide treats 10 acres. Apply as a foliar spray when the wheat is in the 2 to 5 leaf stage.

Fall Seeded Wheat: For control or suppression of the listed weeds, apply 0.8 to 1.2 pints of MONDAK® herbicide (1/2 to 3/4 pound of dicamba plus 1/4 to 3/8 pound of MCPA) per acre. One gallon MONDAK® herbicide treats 7 to 10 acres. Use the higher rates on hard to kill or more mature weeds. Apply as a foliar spray immediately after winter dormancy and before the wheat begins to joint.

NONCROPLAND USE: (Roadsides, fence rows and similar areas.)

For top growth control of field bindweed, Canada thistle, perennial sowthistle, Russian knapweed and leafy spurge, apply 3.2 to 8.4 pints of MONDAK® herbicide (1/2 to 1 pound of dicamba and 1 to 2 pounds of MCPA) per acre. For spot treatment apply 1 teaspoonful of MONDAK® herbicide per 1 to 2 square rods. Apply as a foliar spray to actively growing plants.

GRASS SEED PRODUCTION

(Lawn-type fescues, perennial bentgrasses, perennial ryegrasses, and Kentucky bluegrass.)

FOR USE ONLY IN THE STATES OF IDAHO, OREGON AND WASHINGTON.

Apply to grass stands established at least one year prior to treatment.

For control of:

Corn cockle	Dogfennel	Pepperweed
Cornflower	False dandelion	Pineappleweed
(Bachelor's button)	Field pennycress (farweed)	Prickly lettuce
Corn spurry	Filaree	Red sorrel
Chickweed	Henbit	Shepherdspurse
Crab dock	Knawel	Vetch (common)
Dandelion	Lambsquarters	Wild radish

Apply 1 1/2 pints of MONDAK® herbicide (3 1/4 ounces dicamba acid equivalent plus 7 1/4 ounces MCPA acid equivalent) per acre. Apply to early seeding weeds (4" or less in height) after winter dormancy up to the early boot stage of the grass. For ground application, use a minimum of 20 gallons of total spray per acre.

E.P.A. REG. NO. 876-55-AA
E.P.A. EST. NO. 37507-1AT-01

Net Contents: 30 Gallons

18.0%
2.8%
31.7%
48.5%
100.0%

CORPORATION
Page, Illinois 60611