

### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact flush with plenty of water; for eyes get medical attention. Keep out of reach of children.

### IMPORTANT

Do not apply BANVEL® K herbicide directly to or allow spray drifts to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, sugar beets, beans, or other desirable plants. (Coarse sprays are less likely to drift.)

Do not contaminate domestic or irrigation water by cleaning of equipment or disposal of wastes.

Do not store in direct contact with fertilizer, seed, insecticides, or fungicides.

Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes.

Do not graze livestock on treated areas until 30 days after application. Do not graze treated areas or feed treated forage to dairy cattle prior to harvest.

Do not apply (except as recommended) or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemicals may be washed or moved into contact with their roots.

**BANVEL K SPRAY DRIFT:** Soybeans are sensitive to BANVEL K as it requires only a small amount of chemical per acre for symptoms to appear. Symptoms generally appear about two weeks after drift occurs. Soybeans generally outgrow these initial symptoms without any adverse effect on soybean yields.

To minimize BANVEL K spray drift onto soybeans and other sensitive crops, observe the following precautions:

1. Use coarse sprays.
  - a. Velsicol Chemical Corporation recommends the ACCUTROL™ system as an excellent method for drift reduction. For further information on the ACCUTROL™ system, refer to Velsicol Bulletin A-268.
  - b. When using conventional applications, use 20 gallons or more of water per acre. Keep the spray pressure below 20 p.s.i. For broadcast application use TK5 Floodjet™ tips (or equivalent).
2. Spray with the nozzles as close to the weeds as is practical for good coverage.
3. Do not apply BANVEL K Herbicide to field corn when soybeans are more than 10 inches tall or if the daily temperature is expected to be over 85 F.
4. Make application when there is some air movement but wind is less than 5 miles per hour. If any wind over 5 m.p.h. is blowing in the direction of sensitive crops, do not spray.

### CONTAINER DISPOSAL

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Do not burn empty container.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this product.

# VELSICOL

# BANVEL K

## HERBICIDE

Contains  
1.25 Lbs.  
Dicamba  
and 2.5 Lbs.  
2,4-D Per  
Gallon

ACCUTROL™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

TK5 Floodjet™

### CAUTION

Keep Out of Reach of Children

See Left Panel for Additional Cautions

NET CONTENTS: 5 GALLONS

E.P.A. Reg. No. 876-44-30

## VELSICOL CHEMICAL CORPORATION

341 EAST OHIO STREET, CHICAGO, ILLINOIS 60611

USE

Full season  
(not under  
to legume)

Field use  
Post-emerg

# WARNING

Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water.

# IMPORTANT

Avoid application on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

Avoid contamination of domestic or irrigation waters.

Avoid drift to desirable plants. Do not store near fertilizers, feeds, insecticides or fungicides. Do not apply to newly seeded areas.

# CONTAINER DISPOSAL

Do not reuse empty container. Destroy by burying in safe place. Do not burn.

07102-100



# Active Ingredients:

Dimethylamine Salt of dicamba (3,6-dichloro-g-anisic acid)*	2.41%
Dimethylamine Salts of related acids	0.42%
Inert Ingredients	97.17%
Total	100.00%

\*Equivalent to 2% 3,6-dichloro-g-anisic acid.

# Contents:

# CAUTION

Keep out of the reach of children. See left panel for additional Caution.



VELSICOL CHEMICAL CORPORATION

341 East Ohio Street  
Chicago, Illinois 60611

USDA Reg. No. 876-51

# DIRECTIONS:

For use on lawns and ornamental turf areas for control of clover, chickweed\* (common, mouseear, and perennial or stitchwort), curly dock, dog fennel, knotweed, and sheep sorrel, apply 1 pound of VELSICOL Chickweed-Clover Killer per 500 sq. ft. Granules are to be applied by shaker can. For spot treatment, shake sparingly on areas where control is desired. May be applied in the spring before weeds germinate or as a soil application after weeds are present and actively growing. Make no more than one application per year.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty of any kind, either express or implied, and does not warrant results nor residues no greater than any tolerance which may be established by appropriate governmental agencies, nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, misuse, handling or storage of this material.

ACCEPTED

JUN 22 1971

UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT  
AND ECONOMIC POISON REGISTERED UNDER NO.