

WARNING

May be fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ANTIDOTE

EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

INTERNAL—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

DO NOT REUSE EMPTY CONTAINERS

Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of smoke.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

01151-304AA

VELSICOL®

CHLORDANE 72E.C.

SOIL INSECTICIDE

ACTIVE INGREDIENTS:

Technical Chlordane* 72%

Petroleum Distillate 22%

INERT INGREDIENTS 6%

TOTAL 100%

*Equivalent to 43.2% Octachloro-4, 7-methanotetrahydroindane and 28.8% related compounds. This material contains 8 lbs. Technical Chlordane per gallon.

CONTENTS:

WARNING

KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and Other Precautionary Statements

EPA REGISTRATION NO. 876-55-AA

VELSICOL CHEMICAL CORPORATION, 341 East Ohio St., Chicago, Ill. 60611.

Printed in U.S.A

ACCEPTED
APR 19 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 876-55

DIRECTIONS FOR USE

Do not make foliar applications to corn, vegetable crops.

On corn and vegetable crops, make pre-plant application (where applicable) to soil, after incorporate into soil immediately. Use sufficient coverage of soil surface.

On garden beans, do not graze or feed treated livestock.

For Japanese beetle larvae use on turf — keep treated turf until material has been washed in or rain and grass is dry.

Where a range of rates is given, use the high rate on muck or highly organic soil.

Where row treatment is desirable, application time using suitable row equipment.

A 7-10 inch band is recommended for corn and vegetables. Minimum rates should be based on inch band. For best control, a broadcast application is preferred.

Crop (or Use)	INSECTS
Corn (field & sweet) (use suspended effective August 1, 1976)	White grubs (in the State of Michigan only)
	Cutworms
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peanuts, peas, peppers, pimientos, potatoes, radishes, rutabagas, squash, sweet potatoes, tomatoes, turnips, watermelon.	White grubs (in the State of Michigan only)
	White fringed beetle larva (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)
Federal/State Quarantine Programs (For use on sod and ornamental turf only).	Japanese beetle larvae



WARNING

Contact with skin can cause toxic spray mist. In case of contact with water. Avoid contamination of feed the reach of children.

birds and other wildlife. Birds and bees may be killed. Keep out of any where runoff is likely to occur. Do cleaning of equipment or disposal of only as specified on this label.

OR STORE NEAR HEAT OR OPEN

ANTIDOTE

Illage on skin, wash promptly with r. tablespoonful mustard in tumbler of physician.

NEUTRAL DISPOSAL

EMPTY CONTAINERS

ainers and dispose of wastes by bury- y from water supplies. Containers unching holes in them and burying Keep out of smoke.

Chemical Corporation has no control d conditions of use, which are of ol Chemical Corporation makes no either express or implied, for results y tolerance which may be established al agencies, due to misuse, improper material. Nor does Velsicol Chemical esponsibility for injury to persons, erty arising out of misuse, improper material.

VELSICOL® CHLORDANE 72E.C.

SOIL INSECTICIDE

ACTIVE INGREDIENTS:

Technical Chlordane* 72%

Petroleum Distillate 22%

INERT INGREDIENTS 6%

TOTAL 100%

*Equivalent to 43.2% Octachloro-4, 7-methanotetrahydroindane and 28.8% related com unds. This material contains 8 lbs. Technical Chlordane per gallon.

CONTENTS:

WARNING

KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and Other Precautionary Statements

EPA REGISTRATION NO. 876-55-AA

VELSICOL CHEMICAL CORPORATION, 341 East Ohio

Printed in U S A

ACCEPTED

APR 19 1976

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide registered under EPA Reg. No. 876-55

DIRECTIONS FOR USE

Do not make foliar applications to corn, vegetables, fruit and berry crops.

On corn and vegetable crops, make pre-plant broadcast or row application (where applicable) to soil, after plowing, and disk or incorporate into soil immediately. Use sufficient water for thorough coverage of soil surface.

On garden beans, do not graze or feed treated forage or vines to livestock.

For Japanese beetle larvae use on turf — keep children and pets off treated turf until material has been washed into the soil by watering or rain and grass is dry.

Where a range of rates is given, use the higher rate of Chlordane 72EC on muck or highly organic soil.

Where row treatment is desirable, application should be made at planting time using suitable row equipment.

A 7-10 inch band is recommended for control of white grubs in vegetables. Minimum rates should be based on no less than a 7 inch band. For best control, a broadcast application is recommended.

Crop (or Use)	INSECTS	DOSAGE RATES (Chlordane 72EC/Acre)	
		Broadcast	Row
Corn (field & sweet) (use suspended effective August 1, 1976)	White grubs (in the State of Michigan only)	2.5-4 pints	2 pints
	Cutworms	2.5-4 pints	
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	2.5-5 pints	
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peanuts, peas, peppers, pimentos, potatoes, radishes, rutabagas, squash, sweet potatoes, tomatoes, turnips, watermelon.	White grubs (in the State of Michigan only)	2.5-4 pints	2 pints
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	2.5-5 pints	
Federal/State Quarantine Programs (For use on sod and ornamental turf only).	Japanese beetle larvae	5-10 pints (9-18 ounces per 5,000 square feet)	

