

DANVEL D + 2,4-D
General Weed Control

(Contains 0.23 Pounds dicamba and 0.92 Pounds 2,4-D Per Gallon)

CAUTION:

Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT:

Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present.

Newly seeded lawns may be treated after they have been mowed the second time. To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

Active Ingredients:

Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid)*.....	3.18%
Dimethylamine Salts of related acids.....	.52%
Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid**.....	12.72%
Inert Ingredients.....	83.58%
Total.....	100.00%

* Equivalent to 2.64% 3,6-dichloro-o-anisic acid (dicamba) or 0.23 pounds per gallon.

**Equivalent to 10.57% 2,4-dichlorophenoxyacetic acid or 0.92 pounds per gallon.

CAUTION

Keep Out of the Reach of Children
See left panel for additional cautions

MAY 15 1967
876-64
Velsicol Chemical Corporation
341 East Ohio Street
Chicago, Illinois 60611

Velsicol Chemical Corporation
341 East Ohio Street
Chicago, Illinois 60611

USDA Registration No. _____

DIRECTIONS:

For Use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Teas and Rough. For the control of chicory, clovers, dandelions, docks, dog fennel, English daisy, Frenchweed, knotweed (erect-prostrate), ground ivy, hawkweed, healall, henbit, lambquarter, mallow, morning glory, mustard, oxalis, pennywort, peppergrass, pigweed, plantain (buckhorn, broadleaf, Rugel's), poison ivy, poison oak, Canada thistle, burdock, jimson weed, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, wild astor, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, yarrow and many other broadleaf weeds, apply 6.4 tablespoons (3.2 fluid ounces) of DANVEL D + 2,4-D in 1 to 2 gallons of water per 1000 sq.ft. (Equivalent to .25 Lb. dicamba + 1.0 Lb. 2,4-D amine per Acre). One pint treats 5000 sq.ft. Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (spring and fall) to kill certain hard to kill weeds such as poison ivy, poison oak and English daisy. Weeds that start growing in the fall, such as henbit, may be controlled more effectively with a fall application.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty of any kind, either express or implied, and does not warrant results not to be greater than any tolerance which may be established by appropriate governmental agencies, nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, misuse, handling or storage of this material, whether in accordance with directions or not.

Cameraman's Note
Poor Copy

each of Children
Additional Comments

MAY 15, 1967

THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT
ECONOMIC POISON REGISTER
REGISTER NO. 876-6-10000
SPECIAL COMMENTS

Local Corporation
Street
Illinois 60611

on No. _____

1000 sq.ft. (Equivalent to .25 Lb. dicamba + 1.0 Lb. 2,4-D amine per Acre). One pint treats 5000 sq.ft. Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (spring and fall) to kill certain hard to kill weeds such as poison ivy, poison oak and English daisy. Weeds that start growing in the fall, such as henbit, may be controlled more effectively with a fall application.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty of any kind, either express or implied, and does not warrant results nor assumes no greater than any tolerance which may be established by appropriate governmental agencies, nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, misuse, handling or storage of this material, whether in accordance with directions or not.

Cameraman's Note
Poor Copy