

**VELSICOL**

## AWN & TURF HERBICIDE

(Contains 0.23 lbs. Dicamba and 0.92 lbs. 2,4-D Per Gallon)

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**  
See back panel for additional cautions

E.P.A. REG. NO. 876-64-AA  
E.P.A. EST. NO. 876-TX-2

**Net Contents:**  
**2½ Gallons**

**VELSICOL CHEMICAL CORPORATION**  
**341 E. OHIO ST., CHICAGO, ILL. 60611**

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of dicamba (3,6-dichloro-p-oxalic acid)*	3.18%
Dimethylamine Salts of related acids	.53%
Dimethylamine Salt of 2,4- dichlorophenoxyacetic acid**	12.72%

<b>INERT INGREDIENTS</b>	<b>83.58%</b>
<b>Total</b>	<b>100.00%</b>

\* Equivalent to 2.64% 3,6-dichloro-p-oxalic acid (dicamba) or 0.23 pounds per gallon.

\*\* Equivalent to 10.57% 2,4-dichlorophenoxyacetic acid or 0.92 pounds per gallon.

dicamba  
 acid) ..... 3.18%  
 of related acids .. .52%  
 2,4-  
 to field\*\* ..... 12.72%  
 ..... 83.58%  
 ..... 100.00%  
 1% 3,6-dichloro-p-anisic  
 28 pounds per gallon.  
 7% 2,4-dichlorophenoxy-  
 pounds per gallon.

**CAUTION**

**Hazard to Humans and Domestic Animals.**  
 Caution. Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

**Environmental Hazard.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on label.

**IMPORTANT**

Do not apply on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots.

Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants.

Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water.

Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides.

Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bentgrass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time. To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period.

When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf. Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms

are not acceptable, return at once unopened.

**DIRECTIONS:**

**For Use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Rough.** For the control of burdock, chickweed (mouseear-common-stitchwort), chicory, clover, dandelions, docks, dog fennel, English daisy, Frenchweed, knotweed (erect-prostrate), ground ivy, hawkweed, healall, henbit, jimsonweed, lambsquarters, mallow, morningglory, mustard, oxalis, pennywort, peppergrass, pigweed, plantain (buckhorn, broadleaf, Ruge's), poison ivy, poison oak, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, thistle (Canada), wild aster, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, yarrow and many other broadleaf weeds, apply 6.4 tablespoons (3.2 fluid ounces) of Provel™ in 1 to 2 gallons of water per 1000 sq. ft. (Equivalent to .25 lbs. dicamba + 1.0 lb. 2,4-D amine per Acre.) One pint treats 5000 sq. ft. Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (spring and fall) to kill certain hard-to-kill weeds such as poison ivy, poison oak and English daisy. Weeds that start growing in the fall, such as henbit, may be controlled more effectively with a fall application.

**Limit of Warranty and Liability.** The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances and from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

ACCEPTED

*Sept 9, 1977*

UNDER THE FEDERAL INSECTICIDE  
FUNCTIONALITY ACT OF 1972  
FOR ECHINOCHLORIS ACETIC ACID  
EP UNDER NO. *816-04*

**VELSICOL**

**PROWE**

**LAWN & TURF HERBICIDE**

(Contains 0.23 lbs.  
lbs. 2,4-D Per Gallon)

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
See back panel

E.P.A. REG. NO. 87  
E.P.A. EST. NO. 876

**Net Contents:**  
**2 1/2 Gallons**

**VELSICOL CH**  
**341 E. OHIO**

SICOL

Provan

# & TURF HERBICIDE

(Contains 0.23 lbs. Dicamba and 0.92 lbs. 2,4-D Per Gallon)

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**  
See back panel for additional cautions

E.P.A. REG. NO. 876-64-AA  
E.P.A. EST. NO. 876-TX-2

Net Contents:  
2 1/2 Gallons

VELSICOL CHEMICAL CORPORATION  
841 E. OHIO ST., CHICAGO, ILL. 60611

### ACTIVE INGREDIENTS:

Dimethylamine Salt of dicamba (3,6-dichloro-p-enoic acid)*	3.18%
Dimethylamine Salts of related acids	.88%
Dimethylamine Salt of 2,4- dichlorophenoxyacetic acid**	12.72%

<b>INERT INGREDIENTS</b>	<b>83.22%</b>
<b>Total</b>	<b>100.00%</b>

\* Equivalent to 2.64% 3,6-dichloro-p-enoic acid (dicamba) or 0.23 pounds per gallon.

\*\* Equivalent to 10.57% 2,4-dichlorophenoxyacetic acid or 0.92 pounds per gallon.

CAUTION  
Hazardous  
Caution  
Avoid  
after  
water  
medic  
Environ  
This  
pond  
men  
reco  
MPC  
Do not  
tions  
with  
Av  
on to  
other  
Do  
Do  
plant  
Do  
secti  
Do  
dich  
clove  
after  
smal  
spray  
this  
We  
and  
ing 2  
BRAN

**CAUTION**

Hazard to Humans and Domestic Animals. Caution. Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

**Environmental Hazard.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on label.

**IMPORTANT**

Do not apply on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots.

Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants.

Do not contaminate domestic or irrigation waters.

Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water.

Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides.

Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bentgrass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time. To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period.

When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf. Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms

are not acceptable, return at once unopened.

**DIRECTIONS:**

For Use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Rough. For the control of burdock, chickweed (mouseear-common-stitchwort), chicory, clover, dandelions, docks, dog fennel, English daisy, Frenchweed, knotweed (erect-prostrata), ground ivy, hawkweed, henbit, henbit, limonweed, lambsquarters, mallow, morningglory, mustard, oxalis, pennywort, pepperglass, pigweed, plantain (buckhorn, broadleaf, Ruge's), poison ivy, poison oak, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, thistle (Canada), wild aster, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, yarrow and many other broadleaf weeds, apply 6.4 tablespoons (3.2 fluid ounces) of Provel™ in 1 to 2 gallons of water per 1000 sq. ft. (Equivalent to .25 lbs. dicamba + 1.0 lb. 2,4-D amine per Acre.) One pint treats 5000 sq. ft. Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (spring and fall) to kill certain hard-to-kill weeds such as poison ivy, poison oak and English daisy. Weeds that start growing in the fall, such as henbit, may be controlled more effectively with a fall application.

**Limit of Warranty and Liability.** The exclusive remedy of the user or buyer and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances and from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.