CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Tox. to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Log of apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas freated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product contains as specified on this label.

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water.

Internal—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Form 011..8---130

VELSICOL Belt. 5% Dust

Agricultural Insecticide

ACTIVE INGREDIENTS:

Technical Chlordane*	5.0%
INERT INGREDIENTS	95.0%
TOTAL	100.0%

*Equivalent to 3.0% Octachloro-4,7-methanotetrahydroindane and 2.0% related compounds.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and Additional Precautionary Statements

NET CONTENTS: EPA Reg. No. 876-95-AA

VELSICOL CHEMICAL CORPORATION

341 East Ohio Street, Chicago, Illinois 60611

DIRECTIONS FOR USE

CROP	INSECTS	lb./acre Belt 5% Dust	lb./acre Technical Chlordane	DIRECTIONS
COTTON	Boll weevil	20	. 1	Make application at 4 to .5-day intervals Do not apply rafter bolls open
	Cotton fleahoppers Darkling beetle fall armyworm Tarnished plant bug Thrips	10.30	0515	Apply at cotton treating schedules Do not apply after bolls open
	Grasshoppers	20	1	Do not apply after bolls open.
TOBACCO	Hornworms	20- 4 0	1-2	Repeat as necessary.
CITRUS	Grasshoppers	30-40	1.5-2	Apply to ground cover. Do not apply when fruit is on tree. Do not allow dairy animals to graze in treated groves.

NOTE: Application to greenhouse soil is limited to areas where nonfood crops are grown.

Printed in U.S.A.

