

ACCEPTED
FEB 11 1976
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 876-99

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing of dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children. Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not use this product in any manner not specified on the label.

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water. **Internal**—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes.

IMPORTANT

Do not make foliar applications to corn, vegetables, fruit and berry crops. On field and vegetable crops, make preplant broadcast or row applications (where applicable) to soil after plowing and disk or incorporate into soil immediately.

On berry crops, except strawberries, make application at base of plants and incorporate into soil. Do not apply when fruit is present. Use sufficient water to cover treated area thoroughly. Use on strawberries is for pre-plant treatment only.

On tree fruits make soil application only when no fruit is present.

Where a range of rates is given, use the higher rate of BELT 33.3G on muck or highly organic soil.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

VELSICOL®

Belt® 33.3G

**Agricultural Insecticide Granules
 for Soil Insect Control**

ACTIVE INGREDIENT:

Technical Chlordane* 33.3%

INERT INGREDIENTS 66.7%

TOTAL 100.0%

*Equivalent to 19.98% Octachloro-4,7-methanotetrahydroindane and 13.32% related compounds.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and
 Additional Precautionary Statements

NET CONTENTS: 50 POUNDS

E.P.A. Reg. No. 876-99AA

VELSICOL CHEMICAL CORPORATION

341 East Ohio Street, Chicago, Illinois 60611

DIRECTIONS FOR USE

Crop (or Use)	Insects	Broadcast DOSAGE	
		Pounds of Belt 33.3G Granules/Acre	Lb./acre Technical Chlordane
Corn (field & sweet) (use suspended effective August 1, 1976)	White grubs (in the State of Michigan only)	7.5-12	2.5-4
	Cutworms	7.5-12	2.5-4
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peas, peppers, pimientos, potatoes, radishes, rutabagas, soybeans, squash, sweet potatoes, tomatoes, turnips, watermelon	White grubs (in the State of Michigan only)	7.5-12	2.5-4
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5
Apples, apricots, cherries, nectarines, pears, plums, prunes, quinces, blackberries, grapes, huckleberries, loganberries, raspberries, strawberries	White grubs (in the State of Michigan only)	7.5-12	2.5-4
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5
Strawberries (for pre-plant application only)	Root weevil larvae	30	10
	Rootworms	15-30	5-10
NONCROPLANDS	Imported fire ants*	4.5 (max.)	1.5 (max.)

*For fire ant control make mound application only. Apply thoroughly to mound or feed crops.

NOTE: Where row treatment is desirable, application should be made at planting equipment.

A 7-10 inch band is recommended for control of white grubs in vegetables. Row be in a 7 inch band. Minimum rates should be based on no less than a 7 inch broadcast application is recommended.

BELT 33.3G Applicator Calibration Guide

Applicator	Speed (MPH)	6 lb. Belt 33.3G/Ac. (2 lb. active)	9 lb. Belt 33.3G/Ac. (3 lb. active)	Belt 33.3G/Ac. (1 lb. active)
John Deere	4 MPH	Range 2	Range 2	R
	5 MPH	Notch 3	Notch 11	N
	6 MPH	Range 5	Range 7	R
	8 MPH	Notch 7	Notch 15	N
Gandy	4 MPH	Notch 21	Notch 11	N
	5 MPH	Notch 24	Notch 13	N
	6 MPH	Notch 25	Notch 13	N
	8 MPH	Notch 28	Notch 14	N
Noble Case Oliver	4 MPH	Notch 14	Notch 20	N
	5 MPH	Notch 17	Notch 15	N
	6 MPH	Notch 17	Notch 15	N
New International Harvester	4 MPH	Setting 1.8	Setting 2.4	S
	5 MPH	Setting 2.1	Setting 2.8	S
	6 MPH	Setting 2.4	Setting 3.2	S
A & W Free Flow	4 MPH	Washer K	Washer K	K
	5 MPH	Washer L	Washer M	L
	6 MPH	Washer M	Washer O	M
Atlas Chalmers	Any Speed	Notch 8	Notch 8	N
	4 MPH	Notch 1	Notch 1	N
Winnepolis Moline	4 MPH	Notch 1	Notch 1	N
	5 MPH	Notch 2	Notch 2	N
	6 MPH	Notch 4	Notch 4	N

Using The Chart—Your desired setting is obtained by finding your equipment type in the first vertical column in the chart. Next, select your speed during application in the adjacent column to the right. The recommended setting appears on the same line as your speed, in one of the four columns to the right based on the quantity of Belt 33.3G to be applied per acre. The rates are for 40 inch rows which have 13,069 linear feet of row per acre. If rows of less than 40 inches are used, the linear feet of row per acre will increase and the amount of Belt 33.3G applied will also increase. The following chart indicates the amount of Belt 33.3G per acre that will be applied with various row spacings.

Row Spacing	21
Pounds of Belt 33.3G/Acre	1
	1 1/2
	2

The above settings are applicable to all applicators as they calibrate your own equipment.

ACCEPTED
FEB 11 1976
 Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended for the pesticide registered under EPA Reg. No. 876-99

VELSICOL®

Belt® 33.3G

**Agricultural Insecticide Granules
 for Soil Insect Control**

ACTIVE INGREDIENT:
 Technical Chlordane* 33.3%
INERT INGREDIENTS 66.7%
TOTAL 100.0%

*Equivalent to 19.98% Octachloro-4,7-methanotetrahydroindane and 13.32% related compounds.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and
 Additional Precautionary Statements

NET CONTENTS: 50 POUNDS

E.P.A. Reg. No. 876-99AA

VELSICOL CHEMICAL CORPORATION
 341 East Ohio Street, Chicago, Illinois 60611

DIRECTIONS FOR USE

Crop (or Use)	Insects	DOSAGE RATES			
		Broadcast		Row	
		Pounds of Belt 33.3G Granules/Acre	Lb./acre Technical Chlordane	Pounds of Belt 33.3G Granules/Acre	Lb./acre Technical Chlordane
Corn (field & sweet) (use suspended effective August 1, 1976)	White grubs (in the State of Michigan only)	7.5-12	2.5-4	6	2
	Cutworms	7.5-12	2.5-4		
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5		
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peas, peppers, pimientos, potatoes, radishes, rutabagas, soybeans, squash, sweet potatoes, tomatoes, turnips, watermelon	White grubs (in the State of Michigan only)	7.5-12	2.5-4	6	2
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5		
Apples, apricots, cherries, nectarines, pears, plums, prunes, quinces, blackberries, grapes, huckleberries, loganberries, raspberries, strawberries	White grubs (in the State of Michigan only)	7.5-12	2.5-4		
	White fringed beetle larvae (for use only in 8 S.E. States: AL, FL, GA, LA, MS, NC, SC, TN)	7.5-15	2.5-5		
Strawberries (for pre-plant application only)	Root weevil larvae	30	10		
	Rootworms	15-30	5-10		
NONCROPLANDS	Imported fire ants*	4.5 (max.)	1.5 (max.)		

*For fire ant control make mound application only. Apply thoroughly to mound. Do not contaminate food or feed crops.
NOTE: Where row treatment is desirable, application should be made at planting time using suitable row equipment.
 A 7-10 inch band is recommended for control of white grubs in vegetables. Row treatment of corn should be in a 7 inch band. Minimum rates should be based on no less than a 7 inch band. For best control a broadcast application is recommended.

BELT 33.3G Applicator Calibration Guide

Applicator	Speed (MPH)	6 lb. Belt 33.3G/Ac. (2 lb. active)	9 lb. Belt 33.3G/Ac. (3 lb. active)	12 lb. Belt 33.3G/Ac. (4 lb. active)
John Deere	4 MPH	Range 2	Range 2	Range 2
	5 MPH	Notch 3	Notch 11	Notch 18
	6 MPH	Range 2	Notch 16	Notch 24
	8 MPH	Notch 11	Notch 22	Notch 32
Gandy	4 MPH	Notch 21	Notch 25	Notch 28
	5 MPH	Notch 24	Notch 28	Notch 32
	6 MPH	Notch 25	Notch 30	Notch 35
	8 MPH	Notch 28	Notch 34	
Noble Case Oliver	4 MPH	Notch 14	Notch 20	Notch 27
	5 MPH	Notch 17	Notch 25	Notch 33
	6 MPH	Notch 21	Notch 30	
New International Harvester	4 MPH	Setting 1 B	Setting 2 A	Setting 2 9
	5 MPH	Setting 2 7	Setting 2 B	Setting 3 5
	6 MPH	Setting 2 4	Setting 3 2	Setting 3 8
	8 MPH	Setting 3 0	Setting 4 0	Setting 5 0
A & W Free Flow	4 MPH	Washer 1	Washer K	Washer L
	5 MPH	Washer J	Washer L	Washer N
	6 MPH	Washer I	Washer M	Washer O
	8 MPH	Washer L	Washer O	
Allis Chalmers	Ary Speed	Notch 6	Notch 8	Notch 11
	4 MPH	Notch 1	Notch 3	Notch 6
	5 MPH	Notch 2	Notch 4	
	6 MPH	Notch 3	Notch 5	

Using The Chart—Your desired setting is obtained by finding your equipment type in the first vertical column in the chart. Next, select your speed during application in the adjacent column to the right. The recommended setting appears on the same line as your speed, in one of the four columns to the right, based on the quantity of Belt 33.3G to be applied per acre. The rates are for 40 inch rows which have 13,069 linear feet of row per acre. If rows of less than 40 inches are used, the linear feet of row per acre will increase and the amount of Belt 33.3G applied will also increase. The following chart indicates the amount of Belt 33.3G per acre that will be applied with various row spacings.

Row Spacing	20"	30"	36"	40"
Pounds of Belt 33.3G/Acre	6	8	6.66	6
	12	12	9.99	9
	24	16	13.33	12

The above settings are for typical applicators. Your applicator varies as it gets older. It is best for you to calibrate your equipment using the above as a starting point.

CAUTION
 with skin can cause toxic symptoms. Avoid contact with skin, wash with soap and water and foodstuffs. Keep out of the reach of

out of any body of water. Birds feeding on do not contaminate water by cleaning of es. Do not apply where runoff is likely to manner not specified on the label.

ANTIDOTE
 skin, wash promptly with soap and water. onful of mustard in tumbler of water. Call

INNER DISPOSAL
 Destroy container by burying with waste ke or fumes.

IMPORTANT
 to corn, vegetables, fruit and berry crops. make preplant broadcast or row applica- oil after plowing and disk or incorporate

berries, make application at base of plants not apply when fruit is present. Use suffi- rea thoroughly. Use on strawberries is for ation only when no fruit is present.

an, use the higher rate of BELT 33.3G on

NOTICE
 Corporation has no control over storage, use, which are of critical importance, n makes no representation or warranty, results or residues greater than any toler- ed by appropriate governmental agencies. dling or storage of this material. Nor does n assume any responsibility for injury to or property arising out of misuse, improper material.