

VELSICOL

Banvel 5G

GRANULES

CAUTION

KEEP OUT OF THE
REACH OF CHILDREN

See Below for Additional
Precautionary Statements

E.P.A. Reg. No. 876-103-AA
E.P.A. Est. No. 36716-TN-1

VELSICOL CHEMICAL CORPORATION

341 East Ohio St., Chicago, Illinois 60611

ACCEPTED

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, the pesticide
registered under
EPA Reg. No. 876-103-AA

CAUTION

Avoid contact with skin, eyes and clothing. In case of contact, wash with plenty of water. Do not store near feed or foodstuffs.

STORAGE AND DISPOSAL

- Restrictions:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
- Pesticide Disposal:** Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- Container Disposal:** If plastic container, triple rinse or equivalent. Dispose of plastic or paper containers in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
- General:** Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

IMPORTANT

Do not apply BANVEL granules under desirable trees or other plants, or areas where the roots might extend or where BANVEL might be washed into the root area. Do not contaminate domestic or irrigation waters. Pasture and rangeland grasses may be grazed or harvested for dairy feed (green or dry) 60 days after treatment. There is no waiting period between treatment and grazing for animals other than dairy animals. Do not graze meat animals in treated fields within 30 days of slaughter. Rates in excess of 8.0 lbs. dicamba acid equivalent per acre for spot treatment in noncropland may temporarily stunt or eliminate the growth of desirable grasses in the treated areas.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

07105-130AD

NEW BRUSH AND WEED HERBICIDE

ACTIVE INGREDIENTS:

dicamba (3,6-dichloro-o-anisic acid) 5.0%
Related acids 0.9%

INERT INGREDIENTS: 94.1%
100.0%

NET CONTENTS: 25 POUNDS

DIRECTIONS FOR USE

USE: PASTURE, RANGELAND, AND NONCROPLAND AREAS SUCH AS FENCEROWS, ROADWAYS, WASTELANDS AND SIMILAR AREAS.

WEED/BRUSH	DOSAGE	APPLICATION
Eastern Persimmon	Use 2 level teaspoons of BANVEL 5% Granules per inch diameter of the trunk of the plant (Example: Use 6 teaspoons for a tree with a trunk 3 inches in diameter)	Scatter the granules evenly on the ground within 6 inches of the trunk. Apply BANVEL granules any time after buds start to open and before the leaves and branches stop growing in the summer.
Crotonbush Yarabush	Use 2 heaping tablespoons of BANVEL 5% Granules per 4 feet diameter of canopy	Make application just prior to or in the early part of the rainy season. Scatter the granules uniformly under the canopy of the shrub.
Salt Cedar	Use 100 to 200 pounds of BANVEL 5% Granules per acre (5 to 10 lbs. dicamba acid equivalent)	Make application just prior to or in the early part of the rainy season. Apply BANVEL granules uniformly over the area to be treated.
Canada Thistle Field Bindweed (Morningglory) Russian knawweed Leafy spurge Burr ragweed Skeletonweed Tansy Ragwort	Apply at the rate of 80 to 160 lbs. BANVEL 5% Granules (4 to 8 lbs. dicamba acid equivalent) per acre. For spot treatment apply 0.5 to 1.0 lbs. BANVEL 5% Granules per sq. rod (272 sq. ft.)	For best results, apply BANVEL granules uniformly when plants are actively growing. This would normally be in the spring or fall when plants are putting out new growth.
Bracken fern	Apply at the rate of 120 to 160 lbs. BANVEL 5% Granules (6 to 8 lbs. dicamba acid equivalent) per acre. For spot treatment apply 75 to 100 lbs. BANVEL 5% Granules per sq. rod (272 sq. ft.)	Apply granules uniformly as a pre-emergence application before emergence of the fronds.
Artichoke thistle	Apply at the rate of 20 to 40 lbs. BANVEL 5% Granules (1 to 2 lbs. dicamba acid equivalent) per acre. For spot treatment apply 2 to 4 oz. BANVEL 5% Granules per sq. rod (272 sq. ft.)	Make uniform application of BANVEL granules when plants are actively growing.

BEST DOCUMENT AVAILABLE