

Harmful if swallowed. Avoid contact with the skin. Contact with skin can cause toxic symptoms. Avoid breathing vapors and dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

Keep children or pets off treated ornamental turf until material has been washed into the soil by watering or rain and grass is dry. Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not use in any manner not specified on this label.

CONTAINER DISPOSAL

Do not reuse empty container.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or

residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

DIRECTIONS FOR USE

Make broadcast application. Cover surface uniformly. Treatment should be made in spring or when insects are observed. Do not treat when soil is wet.

For use on Ants, Asiatic Garden Beetle Larvae, Japanese Beetle Larvae, White-fringed Beetle Larvae, White Grubs, Wireworms, and European Chafer Larvae, apply GOLD CREST H-2.5% G just before a rain or use irrigation or sprinklers to wash the material into established turf.

For use on Fire Ants, make broadcast application. Cover surface uniformly. Do not treat when soil is wet. Do not apply to pasture or rangeland.

For other insects, apply GOLD CREST H-2.5% G around the base of trees, under shrubs, along walks, fences and the foundations of buildings.

GOLD CREST H-2.5% G Application Rates for Golf Courses, Parks, Lawns, Ornamental Turf and Cemeteries

INSECT	AMOUNT OF GOLD CREST H-2.5% G PER 5,000 SQ. FT.	LBS. PER ACRE ACTIVE INGREDIENT
Chiggers Earwigs Fleas Ticks	74 oz	1 lb
Ants Asiatic Garden Beetle Larvae European Chafer Larvae Japanese Beetle Larvae White fringed Beetle Larvae White Grubs Wireworms	9 lb	2 lb
Fire Ants	For a single broadcast application 5 3/4 lb	1.25 lb
	For two broadcast applications spaced at 3 to 6 month intervals 2 1/4 3 1/3 lb.	0.50 0.75 lb
	Mound Application Apply 4.8 cups of Gold Crest H-2.5% G (equivalent to 0.56 1.2 ounces of heptachlor) to entrance of nest as well as to an area 3 feet around mound. Treated areas should be examined 3 weeks after treatment and surviving colonies retreated.	

Manufactured and distributed by

V E L S I C O L C H E M I C A L C O R P O R A T I O N