

WARNING

Keep away from human beings, domestic animals, wildlife and pets. If swallowed by human beings, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, the intravenous and oral administration of vitamin K combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of *bis*-hydroxycoumarin.

IN ALL CASES OF HUMAN INGESTION, IMMEDIATELY NOTIFY A PHYSICIAN.

CONTAINER DISPOSAL

Do not reuse empty container. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Form No. 16010-350

VELSICOL

Warfarin

CONCENTRATE

RODENTICIDE

KILLS RATS and MICE

Anti-coagulant for Rodent Control

ACTIVE INGREDIENT:

Warfarin* 0.5%

INERT INGREDIENTS 99.5%

Total 100.0%

*3-(α -acetylbenzyl)-4-hydroxycoumarin

NET CONTENTS: 100 Pounds

WARNING

**Keep Out of the Reach of Children.
See left panel for additional warnings.**

VELSICOL CHEMICAL CORPORATION
341 EAST OHIO ST., CHICAGO, ILL. 60611

IMPORTANT

Place baits in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes.

ACCEPTED

Jul 28 1971

DIRECTIONS

(a) Thoroughly mix one part (by weight) of rodenticide with nineteen (19) parts of dry cereal bait.

(b) Place bait in locations where rats and mice feed, water or travel.

(c) Bait placements along walls, in corners, and in concealed locations where rats and mice feed are desirable.

For Rat Control:

(a) Place at least four ounces of bait at each bait location.

(b) Maintain an uninterrupted supply of bait in the stations for at least 15 days.

For Mouse Control:

(a) Place one tablespoonful of bait at 8-10 foot intervals.

(b) Maintain an uninterrupted supply of bait for at least 15 days.

Note:

If bait molds, decomposes, or becomes sour, replace with fresh materials.

U.S.D.A. Reg. No. 876-115

