



WARNING
Keep away from human beings, domestic animals, wildlife and pets. If swallowed by human beings, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, the intravenous and oral administration of vitamin K combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

IN ALL CASES OF HUMAN INDIGESTION, IMMEDIATELY NOTIFY A PHYSICIAN.

CONTAINER DISPOSAL
Do not reuse empty container. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

NOTICE
Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.
16010-360AA

VELSICOL Warfarin CONCENTRATE

**RODENTICIDE
KILLS RATS and MICE**
Anti-coagulant for Rodent Control
For Commercial Use Only

ACTIVE INGREDIENTS:
Warfarin* 0.5%
INERT INGREDIENTS 99.5%
Total 100.0%

*3-(a-acetonylbenzyl)-4-hydroxycoumarin

NET CONTENTS:

WARNING
Keep Out of the Reach of Children.
See left panel for additional warnings.

E.P.A. Reg. No. 876-115-AA
EPA Est. No. 876-AZ-1
VELSICOL CHEMICAL CORPORATION
341 EAST OHIO ST.
CHICAGO, ILL. 60611

IMPORTANT
Place baits in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes.

- DIRECTIONS**
- (a) Thoroughly mix one part (by weight) of rodenticide with nineteen (19) parts of dry cereal bait.
 - (b) Place bait in locations where rats and mice feed, water or travel.
 - (c) Bait placements along walls, in corners, and in concealed locations where rats and mice feed are desirable.

For Rat Control:
(a) Place at least four ounces of bait at each bait location.
(b) Maintain an uninterrupted supply of bait in the stations for at least 15 days.

For Mouse Control:
(a) Place one tablespoonful of bait at 8-10 foot intervals.
(b) Maintain an uninterrupted supply of bait for at least 15 days.

Note:
If bait molds, decomposes, or becomes sour, replace with fresh materials.

