

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing of dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur.

Do not use this product in any manner not specified on the label.

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water.

Internal—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes.

IMPORTANT

Do not make foliar applications to corn, vegetables, fruit and berry crops. On field and vegetable crops, make preplant broadcast or row applications (where applicable) to soil after plowing and disk or incorporate into soil immediately.

On berry crops, except strawberries, make application at base of plants and incorporate into soil. Do not apply when fruit is present. Use sufficient water to cover treated area thoroughly. Use on strawberries is for pre-plant treatment only.

On tree fruits make soil application only when no fruit is present.

Where a range of rates is given, use the higher rate of BELT 40W.P. on muck or highly organic soil.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

VELSICOL[®]

Belt[®] -40% W.P.

Agricultural Insecticide

ACTIVE INGREDIENT:

Technical Chlordane* 40.0%

INERT INGREDIENTS 60.0%

TOTAL 100.0%

*Equivalent to 24% Octachloro-4,7-methanotetrahydroindane and 16% related compounds

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Left Panel for Antidote and Additional Precautionary Statements

NET CONTENTS: 50 POUNDS

EPA Reg. No. 876-172-AA

VELSICOL CHEMICAL CORPORATION
341 East Ohio Street, Chicago, Illinois 60611

DIRE

CROP (or USE)	
Corn (field & sweet) (use suspended effective August 1, 1976)	White of Mic Cutwo White larvae 8 S.E. GA, LA
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peas, peppers, pimentos, potatoes, radishes, rutabagas, squash, sweet potatoes, tomatoes, turnips, watermelon	White of Mic White larvae 8 S.E. GA, LA
Apples, apricots, cherries, nectarines, pears, plums, prunes, quinces, blackberries, grapes, huckleberries, loganberries, raspberries, strawberries	White of Mic White larvae 8 S.E. GA, LA
Strawberries (for pre-plant application only)	Root Rootw
NONCROPLANDS	Impor

*For fire ant control make mound food or feed crops.

NOTE: Where row treatment is desired use row equipment.

A 7-10 inch band is recommended for broadcast application is recommended