

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Toxic to fish and wildlife. Do not use near any body of water. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. DO NOT USE IN ANY MANNER NOT SPECIFIED ON THIS LABEL.

ANTIDOTE

EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

INTERNAL—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

Treated seeds should be labeled as follows:

POISON TREATED—THIS SEED TREATED WITH CHLORDANE. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.

CONTAINER DISPOSAL

Do not reuse empty container. Rinse equipment and dispose of wastes by burying in noncrop lands away from water supplies. Destroy container by punching holes in it and burying with wastes or by burning. Stay out of smoke and fumes.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling, or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

VELSICOL[®] Belt-40%[®] SEED DRESSING FOR PLANTER BOX USE

ACTIVE INGREDIENT:		
Technical chlordane*	40%	
INERT INGREDIENTS	60%	
	TOTAL	100%

*Equivalent to 24% Octachloro-4, 7-methanotetrahydroindane and 16% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Antidote and
Additional Precautionary Statements

NET CONTENTS: 50 POUNDS

EPA Reg. No. 876-173-AA

VELSICOL CHEMICAL CORPORATION
341 East Ohio Street, Chicago, Illinois 60611

DIRECTIONS FOR USE

BELT[®] 40% Seed Dressing is designed for planter box use at the time of planting for temporary protection of seeds and germinating seedlings from attack by soil insects. Mix directly with seed and use in planters designed to give the correct dosages. Belt[®] 40% Seed Dressing should not be mixed with water or water-containing products. Seeds should be cleaned and well cured before treatment.

Seed	Insects	Dosage BELT 40% Seed Dressing
Corn (field)	Seed corn beetle Seed corn maggot Southern corn rootworm Wireworms	5 ounces (equivalent to 2 ounces technical chlordane) per bushel of seed
Cotton	False wireworm Seed corn maggot Wireworms	
Sorghum	False wireworm Kafir ant Seed corn maggot Wireworms	
Soybeans	Seed corn maggot Wireworms	
Small Grains (Oats, rye, wheat)	False wireworm Seed corn maggot Wireworms	
Rice	False wireworm Seed corn maggot Wireworms	

ACCEPTED
Sept. 28, 1972

876 173

