

CAUTION

Harmful if swallowed. Avoid contact with skin. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water.

Internal—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Manufactured and distributed by

VELSICOL CHEMICAL CORPORATION

DIRECTIONS FOR USE

LOCATION	INSECTS	Amounts of Heptachlor Granule per 1,000 sq. ft.	Amounts of Heptachlor Granule per acre	APPLICATION DIRECTIONS
Lawns, Ornamental Turf, Parks, Golf Courses, Cemeteries, Nurseries	Chinch bug Grasshopper Flea beetle Cricket	7 1/2 oz.	20 lbs.	Apply around the base of trees, under shrubs, along walks, fences and the foundations of buildings.
	Asiatic garden beetle larvae European chafer larvae Japanese beetle larvae White fringed beetle larvae White grubs Wireworms	15 oz.	40 lbs.	Make broadcast application. Cover surface uniformly. Apply just before a rain or use irrigation or sprinkler to wash the material into established turf.
	Fire ants	For a single broadcast application 9 oz.	25 lbs.	BROADCAST SOIL TREATMENT Make broadcast application. Cover surface uniformly. Do not treat when soil is wet. Do not apply to pasture or rangeland.
		For two broadcast applications spaced at 3 to 6 month intervals 3 1/4 - 3 1/2 oz.	10 to 15 lbs.	
Non crop, land and non food crop areas	Fire ants	See application directions	See application directions	MOUND APPLICATION Apply 2.4 cups Heptachlor granular equivalent to 0.56 - 1.2 oz. of heptachlor to entrance of nest as well as to an area 3 feet around mound. Treated areas should be examined 3 weeks after treatment and surviving colonies retreated. Do not graze livestock on treated areas within 1 year of application.

02113 142AC

341 East Ohio Street • Chicago, Illinois 60611