

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.


CONTAINER DISPOSAL

Do not reuse empty container. The empty containers should be rinsed with water at least three times before disposal, and rinse water should be added to spray tank. Discard container in a safe place. Follow State Recommendations regarding container disposal.

IMPORTANT

- Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers and tomatoes, during both the growing and dormant periods. (Coarse sprays are less likely to drift.) Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not allow this product to come in direct contact with seeds, insecticides or fungicides. To avoid injury to desirable plants, equipment used for **BANVEL®-720** should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals.

Do not graze dairy animals on treated areas within 7 days after application.



876-177

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

07252-450 AB

55

© VELSICOL CHEMICAL CORPORATION 1980

VELSICOL BANVEL®



WATER SOLUBLE INDUSTRIAL FOR BRUSH AND BROADLEAF

Contains 1 Pound Dicamba and 1.9** Pound

ACTIVE INGREDIENTS

Dimethylamine salt of dicamba
(3, 6-dichloro-p-anisic acid)* 12.82%
Dimethylamine salts of related acids 3.19%
Dimethylamine salt of 2, 4-dichloro-
phenoxyacetic acid** 24.58%

INERT INGREDIENTS: 59.41%

TOTAL 100.00%

*Equivalent to 10.6% 3, 6-dichloro-p-anisic acid

**Equivalent to 20.4% 2, 4-dichlorophenoxyacetic acid by


AC isomer specific method 6, DOI-5.

CAUTION
KEEP OUT OF REACH
OF CHILDREN

See Left Panel for
Additional Cautions

VE
341

VELSICOL BANVEL®-720



WATER SOLUBLE INDUSTRIAL HERBICIDE FOR BRUSH AND BROADLEAF WEED CONTROL

Contains 1 Pound Dicamba and 1.9** Pounds 2,4-D Per Gallon

ACTIVE INGREDIENTS

Dimethylamine salt of dicamba
(3, 6-dichloro-g-anisic acid)*12.82%
Dimethylamine salts of related acids3.19%
Dimethylamine salt of 2, 4 dichloro-
phenoxyacetic acid**24.58%

INERT INGREDIENTS:59.41%

TOTAL100.00%

*Equivalent to 10.6% 3, 6-dichloro-g-anisic acid
**Equivalent to 20.4% 2, 4-dichlorophenoxyacetic acid by
AC isomer specific method 6, DOI-5.

CAUTION
KEEP OUT OF REACH
OF CHILDREN
See Left Panel for
Additional Cautions

ROLL OR SHAKE DRUM BEFORE USING.

E.P.A. Reg. No. 876-177-AA
E.P.A. Est. No. 876-TX-2
NET CONTENTS:
86

VELSICOL CHEMICAL CORPORATION
341 E. OHIO STREET, CHICAGO, ILLINOIS 60611

USE	
Rights-of-Way (utility, railroad, highway, pipeline) Non-selective forest brush control, fence rows, wasteland and similar noncropland	For the control of brush and trees: Alder Ash Aspen Basswood Cedar Cherry Cottonwood Cucumber tree Gum Guava Dogwood Elm Hemlock Hornbeam Locust Maple Mesquite Oak
For control of annual and deep-rooted perennial broadleaf weeds	Perennial: Canada Cockspur Dandelion Field bindweed Lamb's ears Mullein Annual: Carnation Chickweed Clover Cockle Dandelion Horseweed Lambsquarters Muskgrass
To reduce the risk of injury to persons, read the label carefully.	

... of contact.
... of water for at
... on.

... with water at
... to spray tank.
... ns regarding

... or in locations
... roots. Applica-
... when there is
... flowers, fruit
... and tomatoes,
... e less likely to
... estic purposes.
... insecticides or
... BANVEL®-720
... her agricultural

... oration.


... -177

... ndling, and condi-
... no representation
... nce which may be
... ndling or storage of
... or injury to persons.
... e of this materia.

... CHEMICAL CORPORATION 1980

COL

VEL-720



INDUSTRIAL HERBICIDE BROADLEAF WEED CONTROL

amiba and 1.9** Pounds 2,4-D Per Gallon

CAUTION

KEEP OUT OF REACH
OF CHILDREN

See Left Panel for
Additional Cautions

ROLL OR SHAKE DRUM BEFORE USING.

E.P.A. Reg. No. 876-177-AA

E.P.A. Est. No. 876-TX-2

NET CONTENTS:

VELSICOL CHEMICAL CORPORATION
341 E. OHIO STREET, CHICAGO, ILLINOIS 60611

DIRECTIONS FOR USE

USE	BRUSH WEED	RATE 100 GAL. SPRAY MIX	APPLICATION DIRECTIONS
Rights-of-Way (utility, railroad, highway, pipeline). Non-selective forest brush control, fence rows, wasteland and similar noncropland	For the control of unwanted woody brush including: Alder, Persimmon Ash, Pine Aspen, Poplar Basswood, Sassafras Cedar, Spruce Cherry (cherryblossom) Chinquapin, Serviceberry Cucumber tree, Spicebush Gum, Spruce Guava, Sumac Dogwood, Sycamore Elm, Thornapple Hemlock, Thornberry Hornbeam, Wild plum Locust, Willow Maple, Witchhazel Mesquite, Yucca Oak, and similar woody plant species	1 gallon BANVEL-720 Herbicide plus 99 gallons of water	Hydraulic Spray Application Stem Foliage—High Water Volume Apply 1 BANVEL-720 Herbicide after leaves are fully developed and until three weeks before frost. Treat all stems and foliage to runoff with special emphasis on covering the root crown. Use 200 to 300 gallons of spray mix per acre depending on the height and density of the brush.
		9 gallons BANVEL-720 Herbicide plus 91 gallons of water	Back Pack Mist-Blower Application Stem Foliage—Low Water Volume Apply BANVEL-720 Herbicide after leaves are fully developed and until three weeks before frost. Treat all stems and root crown to runoff. Use mist blower application on brush up to 6 feet tall at the rate of 30 to 35 gallons of spray mix per acre of brush.
		3 gallons BANVEL-720 Herbicide plus 12 gallons of water	Aerial Application 15 Gallons of Spray Mix per Acre Apply BANVEL-720 Herbicide after leaves are fully developed and until three weeks before frost. Apply at the rate of 15 gallons of spray mix per acre of brush.
		3 gallons BANVEL-720 Herbicide plus 27 gallons of water	Aerial Application 30 Gallons of Spray Mix per Acre Apply BANVEL-720 Herbicide after leaves are fully developed and until three weeks before frost. Apply at the rate of 30 gallons of spray mix per acre of brush.
For control of annual and perennial broadleaf weeds	Perennial Broadleaf Weeds and Vines Canada thistle, Perennial ragweed Curly dock, Puncturevine Dalmatian toadflax, Redvine Field bindweed, Russian knapweed (morningglory), Tansy ragwort Leaky spurge, Trevine Milkweed, and many other perennial broadleaf weeds	1 gallon BANVEL-720 Herbicide plus 99 gallons of water	For effective broad spectrum control of annual and perennial broadleaf weeds and vines, apply BANVEL-720 Herbicide when weeds are actively growing. Apply at the rate of 100 gallons of spray mix per acre. If less than 100 gallons of spray mix is applied per acre, increase the concentration of BANVEL-720 Herbicide to maintain the same rate of active ingredients per acre of brush.
	Annual Broadleaf Weeds Carpetweed, Pigweed Chickweed, Purslane Clover, Ragweed Cocklebur, Sheep sorrel Dogfennel, Smartweed English daisy, Velvet leaf Henbit, Wild buckwheat Knapweed, and many other annual broadleaf weeds Lambsquarters Mustard	0.5 gallons BANVEL-720 Herbicide plus 99.5 gallons of water	

To reduce the hazards of spray drift when applying this product by ground or air equipment, apply with a drift reducing agent such as foam spray adjuvant, inert system or with other systems designed to reduce spray drift.