


VELSICOL®

CHLORDANE



**ACTIVE
INGREDIENT:**
 Technical
 Chlordane* 10.0%

**INERT
INGREDIENTS: 90.0%**

Total 100.0%

*Equivalent to 6.0% Octachloro-4, 7-methanotetrahydroindane and 4.0% related compounds.

CAUTION
**KEEP OUT OF
 REACH OF CHILDREN**
 See antidote and additional
 caution statements below.

E.P.A. Reg. No. 876-189-AA

NET CONTENTS 50 lbs.

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children. Keep children or pets off treated turf until material has been washed into the soil by watering or rain and grass is dry.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any

DIRECTIONS FOR USE

Insect	Area Treated	Application Rate Chlordane 10G	Application Directions
Fire Ant	Non-cropland Lawns, ornamental turf, cemeteries, golf courses, parks, industrial sites, roadides, and similar non-cropland	10-15 lbs. (1 to 1 1/2 lbs. technical Chlordane) per acre 1-1 1/2 lbs. per 1,000 square feet	Broadcast Soil Treatment Make broadcast application over surface uniformly. Do not treat when soil is wet. Do not apply to pasture or range land.
	Non-cropland and non-food crop areas	See application directions	Mound Application Apply 1 to 3 cups of Chlordane 10G (equivalent to 0.07 to 0.14 lbs. technical Chlordane) to entrance of nest.

INGREDIENTS: 90.0%

Total 100.0%

Equivalent to 6.0% Octachloro-4, 7-methanotetrahydroindane and 4.0% related compounds.

E.P.A. Reg. No. 876-189-AA

NET CONTENTS 50 lbs.

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children. Keep children or pets off treated turf until material has been washed into the soil by watering or rain and grass is dry.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE

EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

INTERNAL—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

DO NOT REUSE EMPTY CONTAINER

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Manufactured and distributed by

VELSICOL CHEMICAL CORPORATION 341 East Ohio Street • Chicago, Illinois 60611

01110-140AD

Printed in U.S.A.

DIRECTIONS FOR USE

Insect	Area Treated	Application Rate Chlordane 10G	Application Directions
Fire Ant	Non-cropland: Lawns, ornamental turf, cemeteries, golf courses, parks, industrial sites, roadsides, and similar non-cropland.	10-15 lbs. (1-1 1/2 lbs. technical Chlordane) per acre 1-1 1/2-1 3/4 lbs. per 5,000 square feet	Broadcast Soil Treatment Make broadcast application in furrows uniformly. Do not treat when soil is wet. Do not apply to pasture or range land.
	Non-cropland and non-food crop areas	See application directions	Mound Application Apply 2 to 4 cups of Chlordane 10G (equivalent to 0.27-0.54 lbs. technical Chlordane) to entrance of nest as well as to an area 3 feet around mound. Treated areas should be examined 3 weeks after treatment and surviving colonies retreated. Do not graze livestock on treated areas within 1 year of application.

VEGETABLE CROPS

Do not make foliar application to vegetables. Make preplant broadcast or row application (where applicable) to soil after plowing and disk or incorporate into soil immediately. Do not graze or feed forage or vines from treated beans to livestock. Where a range of rates is given, use the higher rate of Chlordane 10G on muck or highly organic soil.

CROP (Or Use)	INSECTS	DOSAGE RATES			
		BROADCAST		ROW (See Note Below)	
		Ounces Chlordane 10G/1,000 sq. ft.	Lb/acre Technical Chlordane	Ounces of Chlordane 10G/1,000 sq. ft.	Lb/acre Technical Chlordane
Beans, beets, broccoli, brussels sprouts, cabbage, cantaloupe, cauliflower, celery, collards, cucumbers, eggplant, kale, kohlrabi, lettuce, melons, okra, onions, peas, peppers, pimentos, potatoes, radishes, rutabagas, squash, sweet potatoes, tomatoes, turnips, watermelon.	White grubs (in the State of Michigan only)	9-14	2-5.4	7	2
	White fringed beetle larvae (for use only in 8 S.E. states: AL, FL, GA, LA, MS, NC, SC, TN)	9-18	2-5.5		

NOTE: Where row treatment is desirable, application should be made at planting time using suitable row equipment. A 7-10 inch band is recommended for control of white grubs in vegetables. Minimum rates should be based on no less than a 7 inch band. For best control, a broadcast application is recommended.