

VELSICOL

ACTIVE INGREDIENTS:

Xylene	63.53%
Heptachlor*	23.29%
Related Compounds	8.18%
INERT INGREDIENTS	5.00%


TOTAL.....100.00%

Heptachlorotetrahydro-4,7-methanoindene

CONTAINS 2 LBS. ACTUAL HEPTACHLOR PER GALLON.

KEEP OUT OF THE REACH
OF CHILDREN

See left panel for Antidote and Additional
Precautionary Statements



E.P.A. Registration No. 876-190-AA

WARNING

May be fatal if swallowed! Do not breathe vapor or spray mist. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed or food-stuffs. Keep out of reach of children. Do not use near heat or open flame. Store this product at temperatures above 50 F.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

Do not apply when weather conditions favor drift from areas treated.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Keep children and pets off treated areas until material is washed into soil or grass is dry.

ANTIDOTE

INTERNAL—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

CONTAINER DISPOSAL

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

0-153 163PR

VELSICOL



NET CONTENTS:

341 EAST OHIO STREET, CHICAGO, ILLINOIS 60611

IMPORTANT

Dilute Heptachlor 2 EC with sufficient water to give adequate coverage. May be applied by air for control of insects in corn and soy beans using 1/2 to 2 gallons of water per acre.

Row treatment for corn should be applied in front of planter packing wheel.

Preplant broadcast application to corn, pineapple, soybeans, snap beans and lettuce should be worked into the soil immediately after treatment.

Do not apply to berries, peppers, and tomatoes after the edible parts start to form.

Repeat applications to turf as necessary to maintain effective control. Keep children and pets off treated area until material is washed into the soil and grass is dry. Do not graze cattle on treated grass.

For control of cabbage maggots use 3/4 pt. per acre and spray in furrow at planting time. May be used as a drench at planting time by using 1/2 pt/50 gal. water. Treating is limited to planting or transplanting only.

For pineapples make preplant soil application only.

Do not make applications to soil within 60 days of harvest.

ACCEPTED

JUN 04 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR THE CONTROL OF PESTS ON REGISTERED
CROPS UNDER THE FEDERAL INSECTICIDE ACT
TO ATTACHED LABELS

Cro
Field C
Corn (F
Soybean
Beans
Lettuce Tomato
Rutaba
Cabbag
Berries Blac blue boys dew rasp
Pineap
Ornam Lawns Golf Co Cemete Nursery

DIRECTIONS FOR USE

IMPORTANT

Dilute Heptachlor 2 EC with sufficient water to give adequate coverage. May be applied by air for control of insects in corn and soy beans using 1/2 to 2 gallons of water per acre.

Row treatment for corn should be applied in front of planter packing wheel.

Preplant broadcast application to corn, pineapple, soybeans, snap beans and lettuce should be worked into the soil immediately after treatment.

Do not apply to berries, peppers, and tomatoes after the edible parts start to form.

Repeat applications to turf as necessary to maintain effective control. Keep children and pets off treated area until material is washed into the soil and grass is dry. Do not graze cattle on treated grass.

For control of cabbage maggots use 3/4 pt. per acre and spray in furrow at planting time. May be used as a drench at planting time by using 1/2 pt/50 gal. water. Treating is limited to planting or transplanting only.

For pineapples make preplant soil application only.

Do not make applications to soil within 60 days of harvest.

Crop (or Use)	Insect	Dosage per Acre Heptachlor 2EC Soil Treatment Only	
		Row 10"-14"	Broadcast
Field Crops			
Corn (Field)	Seed Corn beetle	3 pts	3 qts.
	Corn billbug		4 1/2 qts.
	Corn rootworms, seed corn maggot, rough headed corn stalk borer	2 1/4-4 1/2 pts	2 1/4-4 1/2 qts.
	Ants, cutworms, green June beetle larvae, Japanese beetle larvae, white grubs, white-fringed beetle larvae, grape colaspis	3 3/4-6 pts.	3 3/4-6 qts.
		Broadcast Soil Treatment Only	
Soybeans	Grape colaspis, Japanese beetle larvae, seed corn maggot, white grubs, wireworms	3 3/4-6 qts	
Beans (snap)	Wireworms	4 1/2 qts.	
Lettuce, Peppers, Tomatoes	Wireworms	6 qts.	
Rutabagas	Root maggots	6 9/4 qts. Preplant	
Cabbage	Cabbage maggot	3/4 pt.	
Berries Blackberries, blueberries, boysenberries, dewberries, raspberries	Strawberry root weevil, Strawberry root maggot, White grubs	8 1/4 qts.	
Pineapple	Anomala beetle	6 qts. Soil	
		Dosage	
		1000 sq ft	Acre
Ornamental Turf, Lawns, Golf Courses, Cemeteries, Nurseries	Ants, Asiatic garden beetle, European chafer larvae, Japanese beetle larvae, white fringed beetle larvae, white grubs, wireworms	3 oz	4 1/2 qts
	Chiggers, earwigs, fleas, ticks	1 oz	2 1/2 qts

ACCEPTED

JUN 04 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC INDIAN REG. SUPP-
ED UNDER N.C. 111-1. SUBJECT
TO ATTACHED COMMENTS

WARNING

May be fatal if swallowed. Do not breathe dust. Do not get on skin. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water.

Internal—Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

DO NOT REUSE EMPTY CONTAINER.

Destroy container by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation of warranty, either express or implied for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Manufactured and distributed by

VELSICOL CHEMICAL CORPORATION

341 East Ohio S

02111 141AA

LOCATION	INSECTS	Amount Heptachlor Granules 1,000 s
Lawns, Ornamental Turf, Parks, Golf Courses, Cemeteries, Nurseries	Chiggers, Earwigs, Fleas, Ticks	3 3/4
	Ants, Asiatic garden beetle larvae, European chafer larvae, Japanese beetle larvae, White fringed beetle larvae, White grubs, Wireworms	7 1/2
	Fire ants	For single broad application 4 1/2 For broad application space 3 to 6 inches 1 3/4 to 2
Non-cropland and non food crop areas	Fire ants	See application directions