

WARNING

Harmful if swallowed, inhaled or absorbed through the skin or eyes. Do not get in eyes, on skin or on clothing. Avoid breathing dust or spray mist. Keep out of the reach of children. Avoid contamination of feed and foodstuff.

In case of skin contact, wash immediately with soap and water. For eyes flush with plenty of water for at least 15 minutes and get medical attention.

Wear natural rubber gloves and protective clothing when handling. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Observe all precautions on labels of all products used in tank mixtures.

DO NOT USE IN ANY MANNER NOT SPECIFIED ON THIS LABEL.

IMPORTANT

Prevent drift of product to desirable plants. Do not use equipment used for PROBE 75 for any other purpose before thoroughly cleaning. Do not store in direct contact with fertilizers, seeds, insecticides or fungicides.

Do not graze or feed forage from treated areas to livestock. Do not rotate with any other crops within one year of treatment, except cover crops which must be plowed under and not grazed.

When applied according to label directions and under normal growing conditions, PROBE 75 WP will not harm the treated crop. Over-application may result in stunting, crop injury or soil residue. Uneven application of PROBE 75 WP can result in erratic weed control, or crop injury.

Seeding disease, cold weather, deep planting, excessive moisture, high salt concentration, or drought may weaken crop seedlings and increase possibility of damage from PROBE 75 WP. Under these conditions, delayed crop development or reduced yields may result.

CONTAINER DISPOSAL

Do not reuse or burn empty containers. Dispose of empty containers by burying in a safe place away from water supplies.

NOTICE

Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of vital importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling, or change of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons or property, or damage to crops, resulting from the use of this material, when the same is used in accordance with the label directions.

VELSICOL

**Probe
75**

**WETTABLE
POWDER**

Pre- And
Post-Emergence
Selective
Herbicide

ACTIVE INGREDIENTS:

Methazole* 75.0%

Related Compounds 3.9%

INERT INGREDIENTS 21.1%

Total 100.0%

* 1,1-dimethyl-2-methyl-4-methyl-2,4-dinitrophenol

WARNING

**KEEP OUT OF
REACH OF CHILDREN**

See Side Panel for
Additional Precautions

NET CONTENTS: 4 LBS.

E.P.A. Reg. No. 876-105

VELSICOL CHEMICAL CORPORATION
341 EAST OHIO STREET, CHICAGO, ILLINOIS 60611

DIRECTIONS FOR USE

This material is designed to be used either as a preemergence or postemergence application. In cotton production, PROBE 75 WP may be applied preemergence at 3 to 4 lbs. per acre or postemergence at 2 to 3 lbs. per acre. In other states, make no more than 4 applications per season. In other states, make no more than 4 applications per season.

PREEMERGENCE

STATE	WEED	DOSAGE PROBE 75/ACRE	APPLICATION DIRECTIONS
Alabama	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Arkansas	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Georgia	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Louisiana	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Mississippi	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Missouri	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
New Mexico	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
North Carolina	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Oklahoma	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
South Carolina	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Tennessee	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.
Texas	Handgrass	1.5 to 2.0 lbs.	Make application 2 to 4 weeks before planting.

POSTEMERGENCE — PROBE ALONE

STATE	WEED	DOSAGE PROBE 75/ACRE	APPLICATION DIRECTIONS
Alabama	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Arkansas	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Georgia	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Louisiana	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Mississippi	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Missouri	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
New Mexico	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
North Carolina	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Oklahoma	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
South Carolina	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Tennessee	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Texas	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.

POSTEMERGENCE — TANK MIX WITH MSMA

STATE	WEED	DOSAGE PROBE 75/ACRE	APPLICATION DIRECTIONS
Alabama	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Arkansas	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Georgia	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Louisiana	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Mississippi	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Missouri	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
New Mexico	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
North Carolina	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Oklahoma	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
South Carolina	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Tennessee	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.
Texas	Handgrass	1.5 to 2.0 lbs.	Apply only as a directed spray to the base of cotton plants.

MIXING DIRECTIONS

For preemergence application, mix 1.5 to 2.0 lbs. of PROBE 75 WP with 20 to 40 gallons of water per acre. For postemergence application, mix 1.5 to 2.0 lbs. of PROBE 75 WP with 20 to 40 gallons of water per acre. In Texas, Oklahoma and New Mexico, if a preemergence application is made, make only one postemergence application per season. If no preemergence application has been made, two postemergence applications may be made per season.