



GOLD CREST

PROMARTM meal

**RODENTICIDE FOR USE ONLY BY PROFESSIONAL APPLICATORS
for Control of Commensal Mice and Rats**

CAUTION

**KEEP OUT OF THE
REACH OF CHILDREN**

**SEE BELOW FOR
ADDITIONAL CAUTIONS**


ACTIVE INGREDIENT:

Diphacinone
(2-Diphenylacetyl-1,
3-Indandione) 0.005%

INERT INGREDIENTS: 99.995%

TOTAL 100.000%

E.P.A. Reg. No. 876-197-AA



CAUTION

Keep out of the reach of children, pets, wildlife, and domestic animals.

NET CONTENTS:

DIRECTIONS FOR USE

Place baits in



GOLD CREST

PROMARTM meal

**RODENTICIDE FOR USE ONLY BY PROFESSIONAL APPLICATORS
for Control of Commensal Mice and Rats**

CAUTION

**KEEP OUT OF THE
REACH OF CHILDREN**


SEE BELOW FOR
ADDITIONAL CAUTIONS

ACTIVE INGREDIENT:

Diphacinone
(2-Diphenylacetyl-1,
3-Indandione) 0.005%

INERT INGREDIENTS: 99.995%
TOTAL 100.000%

E.P.A. Reg. No. 876-197-AA



CAUTION

Keep out of the reach of children, pets,
wildlife, and domestic animals.

NET CONTENTS:

DIRECTIONS FOR USE



CAUTION

Keep out of the reach of children, pets, wildlife, and domestic animals.

GOLD CREST PROMAR Meal should be placed in tamperproof bait boxes or in areas which are inaccessible to children, pets, wildlife, and domestic animals.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol). IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling, or storage of this material.

NET CONTENTS:

DIRECTIONS FOR USE

Place baits in tamperproof bait stations along walls, in corners, and in concealed locations where rats and mice can feed undisturbed.

Bait placements should be checked daily to make sure that an uninterrupted supply of bait is available. Maintain an uninterrupted supply of bait for at least 15 days. Continue until all signs of feeding have stopped.

RATS: Place 4 to 16 ounces of bait per placement.

MICE: Tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ ounce) of bait should be placed at 8 to 12 foot intervals.

Bait placements should be kept permanently to prevent new infestation by migratory rats and mice.

For best results use only fresh bait. Destroy bait that gets mouldy, sour, dirty or infested with insects.

VELSICOL CHEMICAL CORPORATION

341 East Ohio Street • Chicago, Illinois 60611