

**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

**CONTAINER DISPOSAL**

Do not reuse empty container. The empty containers should be rinsed with water at least three times before disposal, and the rinse water should be added to spray tank. Discard container in a safe place. Do not burn. Follow State Recommendations regarding container disposal.

**IMPORTANT**

Do not use around the home or recreation areas.

Consult your State authorities for possible aerial application or other application restrictions. Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, and tomatoes, during both the growing and dormant periods. (Coarse sprays are less likely to drift.) Do not contaminate irrigation ditches or water used for domestic purposes. Do not allow this product to come in direct contact with fertilizers, seeds, insecticides or fungicides. Consult your State Experimental Station or Extension Service Weed Specialists. To avoid injury to desirable plants, equipment used for Brush Buster III should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals.

When using 1 gallon of Brush Buster III per acre, do not graze dairy animals on treated areas within 6 weeks after application.

When using 3 gallons of Brush Buster III per acre, do not graze dairy animals on treated areas within 60 days after application.

Do not graze meat animals on treated areas within 2 weeks of slaughter.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling or storage of this material.


01274 100AC

VELSICOL CHEMICAL CORPORATION

**VELSICOL®**

# Brush Buster

**HERBICIDE**



**KEEP OUT OF THE REACH OF CHILDREN CAUTION**

Contains 1 pound of 1 pound 2,4-D and 2, 4, 5-T per Gallon

**ACTIVE INGREDIENTS:**  
Dimethylamine salt of dicamba (3, 6-dichloro-p-anisic acid) .....  
Dimethylamine salts of related acids .....  
Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid\* .....  
Dimethylamine salt of 2, 4-trichlorophenoxyacetic acid\*\* .....

**INERT INGREDIENTS:**  
TOTAL .....  
\*Equivalent to 10.70%  
\*\*Equivalent to 10.70%  
\*\*\*Equivalent to 10.70%

**See Left Panel for Additional Cautions**

E.P.A. Reg. No. 876-206-00  
E.P.A. Est. No. 876-TX-2

**NET CONTENTS: 5 Gallons**

**ACCEPTED**  
5 MAY 1980  
876-207

**SHAKE WELL**

**VELSICOL CHEMICAL CORPORATION • 341 East Ohio Street**

75

# Brush Buster™ III

HERBICIDE

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use on firetrails and lanes; forest lands, management areas, plantations, and stumplands; rights-of-way; highways, pipelines, powerlines, utilities, roadsides, roadways, etc.; and pastures; or for any other use not included on this label.

Areas specified on this label can be treated ONLY if they are NOT included as part of those areas mentioned in the above paragraph.

For nonselective brush and broadleaf weed control along fencerows, wastelands and noncropland. Do not use around the home or recreation areas.

1 pound Dicamba,  
2,4-D and 1 pound  
per Gallon

**INGREDIENTS:**

|                      |         |
|----------------------|---------|
| the salt of dicamba* |         |
| o-anisic             | 12.78%  |
| salts of             | 1.72%   |
| the salt of 2, 4-    |         |
| oxyacetic            | 12.78%  |
| the salt of 2, 4, 5- |         |
| oxyacetic            | 12.08%  |
| INGREDIENTS:         | 60.04%  |
| TOTAL                | 100.00% |

\*Equivalent to 10.70% 3, 6-dichloro-p-anisic acid.  
Equivalent to 10.70% 2, 4-dichlorophenoxyacetic acid.  
Equivalent to 10.70% 2, 4, 5-trichlorophenoxyacetic acid.

| BRUSH OR WEED  | AMOUNTS FOR PREPARATION OF SPRAY MIX   | APPLICATION DIRECTIONS   |
|--|--|--|
| <b>Perennial Broadleaf Weeds and Vines</b><br>Canada thistle<br>Curly dock<br>Dalmatian toadflax<br>Field bindweed (morningglory)<br>Leafy spurge<br>Milkweed<br>and many other perennial broadleaf weeds                            | Perennial ragweed<br>Puncturevine<br>Redvine<br>Russian knapweed<br>Tansy ragwort<br>Tansy   | Add 1/2 gallon of Brush Buster III Herbicide (equivalent to 1/2 lb dicamba plus 1/2 lb 2, 4-D plus 1/2 lb 2, 4, 5-T acid) to 49 1/2 gallons of water to make 50 gallons of spray mix.<br><br>For effective broad spectrum control of annual and perennial broadleaf weeds and vines, apply Brush Buster III Herbicide when weeds are actively growing.<br><br>Apply at the rate of 50 gallons of spray mix per 1/2 acre, (about 22,000 square feet).<br><br>If less than 50 gallons is applied per 1/2 acre, increase the concentration of Brush Buster III Herbicide to maintain the same rate of active ingredient per 1/2 acre.   |
| <b>Annual Broadleaf Weeds</b><br>Carpetweed<br>Chickweed<br>Clover<br>Cocklebur<br>Dogfennel<br>English daisy<br>Herb<br>Knawel<br>Lambsquarters<br>and many other annual broadleaf weeds  | Mustard<br>Pigweed<br>Purslane<br>Ragweed<br>Sheep sorrel (red sorrel)<br>Smartweed<br>Vetch<br>Wild buckwheat   | Add 1 quart of Brush Buster III Herbicide (equivalent to 1/2 lb dicamba plus 1/2 lb 2, 4-D plus 1/2 lb 2, 4, 5-T acid) to 49 1/2 gallons of water to make 50 gallons of spray mix.<br><br>For effective broad spectrum control of annual and perennial broadleaf weeds and vines, apply Brush Buster III Herbicide when weeds are actively growing.<br><br>Apply at the rate of 50 gallons of spray mix per 1/2 acre, (about 22,000 square feet).<br><br>If less than 50 gallons is applied per 1/2 acre, increase the concentration of Brush Buster III Herbicide to maintain the same rate of active ingredient per 1/2 acre.<br><br><b>FOR SPOT TREATMENT</b><br>Mix 1 to 2 ounces of Brush Buster III Herbicide in 2 to 3 gallons of water to treat 700 square feet. |
| <b>STEM FOLIAGE APPLICATION—HIGH WATER VOLUME</b>  |  |  |
| <b>Woody Brush</b><br>Alder<br>Ash<br>Aspen<br>Basswood<br>Cedar<br>Chinquapin<br>Cherry<br>Cucumber tree<br>Gum<br>Guava<br>Dogwood<br>Elm<br>Hemlock<br>Hornbeam<br>Locust<br>Maple<br>Mesquite<br>and similar woody plant species | Oak<br>Persimmon<br>Pine<br>Poplar<br>Sassafras<br>Schinus (Christmasberry)<br>Serviceberry<br>Spicebush<br>Spruce<br>Sumac<br>Sycamore<br>Thornapple<br>Thornberry<br>Wild plum<br>Willow<br>Witchhazel<br>Yaupon | Add 1/2 gallon Brush Buster III Herbicide (equivalent to 1/2 lb dicamba plus 1/2 lb 2, 4-D plus 1/2 lb 2, 4, 5-T acid) to 49 1/2 gallons of water to make 50 gallons of spray mix.<br><br>Apply Brush Buster III Herbicide after leaves are fully developed and until three weeks before frost.<br><br>Treat all stems and foliage to runoff with special emphasis on covering the root crown. Apply at the rate of 50 to 75 gallons of spray mix per 1/2 acre of brush depending on the height and density of the brush.  |
| <b>AERIAL APPLICATION*—15 GALLONS OF SPRAY MIX/ACRE</b>  |  |  |
|  | Add 3 gallons of Brush Buster III Herbicide (equivalent to 3 lbs dicamba plus 3 lbs 2, 4-D plus 3 lbs 2, 4, 5-T acid) to 27 gallons of water to make 30 gallons of spray mix.                                      | Apply Brush Buster III Herbicide after leaves are fully developed and until three weeks before frost.<br><br>Apply at the rate of 15 gallons of spray mix per acre of brush.   |
| <b>AERIAL APPLICATION*—30 GALLONS OF SPRAY MIX/ACRE</b>  |  |  |
|  | Add 3 gallons of Brush Buster III Herbicide (equivalent to 3 lbs dicamba plus 3 lbs 2, 4-D plus 3 lbs 2, 4, 5-T acid) to 27 gallons of water to make 30 gallons of spray mix.                                      | Apply Brush Buster III Herbicide after leaves are fully developed and until three weeks before frost.<br><br>Apply at the rate of 30 gallons of spray mix per acre of brush.   |

\*Before making aerial applications, consult your State or Local authorities for possible aerial application restrictions. To reduce the hazards of spray drift when applying this product by ground or air equipment, apply with a drift reducing agent such as foam spray adjuvant, inert system or with other systems designed to reduce spray drift.

BE WELL BEFORE USING

110 Street, Chicago, Illinois 60611

76

77