

ACCEPTED

Oct 5, 1977

CAUTION

Do not apply BANVEL X.P. pellets under desirable trees or other plants, or areas where their roots might extend. Do not contaminate domestic or irrigation waters. Pasture and rangeland grasses may be grazed or harvested for dairy feed (green or dry) 60 days after treatment. There is no waiting period between treatment and grazing for animals other than dairy animals. Do not graze meat animals in treated fields within 30 days of slaughter. To avoid injury to desirable plants, equipment used for BANVEL X.P. pellets should be thoroughly cleaned before reuse to handle or apply any other agricultural chemical. The dosage required for control of these brush-type plants will temporarily stunt or eliminate the growth of desirable grasses in the treated areas.

**IMPORTANT**

Do not apply BANVEL X.P. pellets under desirable trees or other plants, or areas where their roots might extend. Do not contaminate domestic or irrigation waters. Pasture and rangeland grasses may be grazed or harvested for dairy feed (green or dry) 60 days after treatment. There is no waiting period between treatment and grazing for animals other than dairy animals. Do not graze meat animals in treated fields within 30 days of slaughter. To avoid injury to desirable plants, equipment used for BANVEL X.P. pellets should be thoroughly cleaned before reuse to handle or apply any other agricultural chemical. The dosage required for control of these brush-type plants will temporarily stunt or eliminate the growth of desirable grasses in the treated areas.

**CONTAINER DISPOSAL**

Rinse equipment and container and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not burn.

**DIRECTIONS FOR USE**

USE	BRUSH/TREE	DOSAGE	APPLICATION
Pastures, rangeland and non-crop land areas, such as fence rows, high ways, utilities, railroads, and pipeline right of way wash and fill areas.	Eastern Persimmon	Use 1 level teaspoonful of BANVEL X.P. pellets per inch diameter of trunk of the plant. (For example, use 3 teaspoonfuls for a tree with a trunk 3 inches in diameter.)	Scatter the pellets evenly on the ground within 6 inches of the trunk.  Apply BANVEL X.P. pellets any time after buds start to open and before the leaves and branches start growing in the summer.
	Creosotebush Tartar	Use 1 heaping teaspoonful of BANVEL X.P. pellets per 4 feet diameter of shrub.	Make application just prior to or in the early part of the rainy season. Scatter the pellets uniformly under the canopy of the shrub.
	Soft Cedar	Use 50 to 100 pounds of BANVEL X.P. pellets per acre (or 10 lbs. per 100 sq. ft.) and equivalent.	Make application just prior to or in the early part of the rainy season. Apply BANVEL X.P. pellets uniformly over the area to be treated.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse or improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of the use, application, handling or storage of this material.

**VELSICOL**

**BANVEL X.P.**

**BRUSH AND TREE**

**Easy to Use**

ACTIVE INGREDIENT

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane

95.0%

NET WEIGHT

EPA Reg. No. 876-178 AA

NET WEIGHT 50 POUNDS


**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

See Side Panel for Additional Cautions

**VELSICOL CHEMICAL CORP.**

341 E. OHIO STREET, CHICAGO, ILLINOIS 60611



May 1968 - 10/1/68

**IMPORTANT**

Do not apply **BANVEL X.P.** pellets under desirable trees or other plants, or areas where their roots might extend, or where **BANVEL X.P.** pellets might be washed into the root area. Do not contaminate domestic or irrigation waters. Pasture and rangeland grasses may be grazed or harvested for dairy feed (green or dry) 60 days after treatment. There is no waiting period between treatment and grazing for animals other than dairy animals. Do not graze meat animals in treated fields within 30 days of slaughter. To avoid injury to desirable plants, equipment used for **BANVEL X.P.** pellets should be thoroughly cleaned before reuse to handle or apply any other agricultural chemical. The dosage required for control of these brush-type plants will temporarily stunt or eliminate the growth of desirable grasses in the treated areas.

**CONTAINER DISPOSAL**

Rinse equipment and container and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not burn.

**DIRECTIONS FOR USE**

U.F.	BRUSH/TREE	DOSAGE	APPLICATION
1.0	Brush (permanently)	Use 1 pound per 100 sq ft of <b>BANVEL X.P.</b> pellets per acre (1000 sq ft) of the trunk of the plant. The application should be made in a band 6 to 8 inches wide around the trunk.	1. Scatter the pellets evenly on the ground within 6 inches of the trunk. 2. Apply <b>BANVEL X.P.</b> pellets to the ground around the trunk of the plant. Do not apply to the trunk of the plant.
1.0	Tree (permanently)	Use 1 pound per 100 sq ft of <b>BANVEL X.P.</b> pellets per acre (1000 sq ft) of the trunk of the plant. The application should be made in a band 6 to 8 inches wide around the trunk.	1. Scatter the pellets evenly on the ground within 6 inches of the trunk. 2. Apply <b>BANVEL X.P.</b> pellets to the ground around the trunk of the plant. Do not apply to the trunk of the plant.
1.0	Tree (temporary)	Use 1 pound per 100 sq ft of <b>BANVEL X.P.</b> pellets per acre (1000 sq ft) of the trunk of the plant. The application should be made in a band 6 to 8 inches wide around the trunk.	1. Scatter the pellets evenly on the ground within 6 inches of the trunk. 2. Apply <b>BANVEL X.P.</b> pellets to the ground around the trunk of the plant. Do not apply to the trunk of the plant.

**NOTICE** - This product is a restricted use pesticide. It is not to be used by anyone who is not registered with the EPA as a pesticide applicator. For more information on this and other restricted use pesticides, see the label.

# Banvel X.P.

## PELLETS

FOR CONTROL OF UNDECIDABLE

BRUSH AND TREES

**Easy to Use**

**ACTIVE INGREDIENTS**

Dicamba (3,6-dichloro-o-anisic acid) 10.0%  
Related acids 1.8%

**INERT INGREDIENTS**

88.2%  
TOTAL 100.0%

E.P.A. Reg. No. 876 178 AA

NET WEIGHT, 50 POUNDS


### CAUTION

**KEEP OUT OF REACH OF CHILDREN**

See Side Panel for Additional Cautions

**VELSICOL CHEMICAL CORP.**

341 E. OHIO STREET, CHICAGO, ILLINOIS 60611



**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

**CONTAINER DISPOSAL**

Do not reuse empty container. The empty containers should be rinsed with water at least three times before disposal, and the rinse water should be added to spray tank. Discard container in a safe place. Do not burn. Follow label recommendations regarding container disposal.

**IMPORTANT**

Do not apply (except as recommended) on or near desirable plants, or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers and tomatoes, during both the growing and dormant periods. (Coarse sprays are less likely to drift.) Do not contaminate irrigation ditches or water used for domestic purposes. Do not allow this product to come in direct contact with seeds, insecticides or fungicides. To avoid injury to desirable plants, equipment used for Brush Buster should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals.

When using 1 gallon of Brush Buster per acre, do not graze dairy animals on treated areas within 7 days after application.

When using 3 gallons of Brush Buster per acre, do not graze dairy animals on treated areas within 60 days after application.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

07262-413AA

**VELSICOL®**

**Brush Buster**

**HERBICIDE**

**CAUTION  
KEEP OUT OF THE  
REACH OF CHILDREN**

See left panel for  
additional cautions

To  
age

#### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

#### CONTAINER DISPOSAL

Do not reuse empty container. The empty containers should be rinsed with water at least three times before disposal, and the rinse water should be added to spray tank. Discard container in a safe place. Do not burn. Follow State Recommendations regarding container disposal.

#### IMPORTANT

Do not apply (except as recommended) on or near desirable plants, or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers and tomatoes, during both the growing and dormant periods. (Coarse sprays are less likely to drift.) Do not contaminate irrigation ditches or water used for domestic purposes. Do not allow this product to come in direct contact with seeds, insecticides or fungicides. To avoid injury to desirable plants, equipment used for Brush Buster should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals.

When using 1 gallon of Brush Buster per acre, do not graze dairy animals on treated areas within 7 days after application.

When using 3 gallons of Brush Buster per acre, do not graze dairy animals on treated areas within 60 days after application.

**NOTICE:** Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

07262-403AA

# VELSICOL

# Brush Buster

## HERBICIDE

### CAUTION

KEEP OUT OF THE  
REACH OF CHILDREN

See left panel for  
additional cautions

**VELSICOL<sup>®</sup>**

**Brush Buster**

**HERBICIDE**

**CAUTION  
KEEP OUT OF THE  
REACH OF CHILDREN**

**See left panel for  
additional cautions**

th  
be

in  
air  
out  
ch  
py-  
ls.  
es  
in  
ir-  
ed

als

als

ng.  
on  
es  
al  
col  
oil