


VELSICOL

# Brush Buster™

HERBICIDE



Contains 1 Pound Dicamba,  
1.9 Pounds\*\* 2,4-D Per Gallon.

**KEEP OUT OF  
THE REACH OF  
CHILDREN**

## CAUTION

See Left Panel for  
Additional Cautions

E.P.A. Reg. No. 876-205-AA  
E.P.A. Est. No. 876-TX-2

NET CONTENTS: 2½ Gallon

### ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro- <i>o</i> -anisic acid)*	12.82%
Dimethylamine salts of related acids	3.19%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid**	24.58%

INERT INGREDIENTS	59.41%
TOTAL	100.0 %

\*Equivalent to 10.6% 3,6-dichloro-*o*-anisic acid  
(dicamba), or 1 lb./gallon.

\*\*Equivalent to 20.4% 2,4-dichlorophenoxyacetic  
acid by isomer specific AOAC method 6.275-6.279,  
13th edition, or 1.9 lbs./gallon.

**SHAKE WELL BEFORE USING**

VELSICOL CHEMICAL CORPORATION • 341 East Ohio Street, Chicago, Illinois 60611

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND TO DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**

Do not apply (except as recommended) on or near desirable plants, or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers and tomatoes, during both the growing and dormant periods. (Coarse sprays are less likely to drift.) Do not contaminate irrigation ditches or water used for domestic purposes. Do not allow this product to come in direct contact with seeds, insecticides or fungicides. To avoid injury to desirable plants, equipment used for Brush Buster should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals. When using 1 gallon of Brush Buster per acre, do not graze dairy animals on treated areas within 7 days after application.

When using 3 gallons of Brush Buster per acre, do not graze dairy animals on treated areas within 60 days after application.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL.** Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under Subtitle C of the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL**

**Metal**

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

**Plastic**

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration if permitted by State and local authorities.

**LIMITED WARRANTY AND LIABILITY**

**NOTICE:** Read the Limited Warranty and Liability before buying or using this product. If terms are not acceptable, return it at once unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and the Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions or from mixing this product before use with any substance except as recommended by the product's label, (ii) arising from handling or storage in violation of label instructions, (iii) for all indirect, special or consequential damages, (iv) when not reported to the company within one year of discovery, and (v) arising from product not used within the label-designated shelf life or four years from the date of purchase, whichever occurs first. **THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For nonselective brush and broadleaf weed control along fencerows, drainage ditch banks, highway rights-of-way, waste-land and similar noncropland.

BRUSH OR WEED	AMOUNTS FOR PREPARATION OF SPRAY MIX	APPLICATION DIRECTIONS
<b>Perennial Broadleaf Weeds and Vines</b> Canada thistle      Leaky spurge      Redvine Curly dock          Milkweed            Russian knapweed Dalmatian toadflax      Perennial ragweed      Tansy ragwort Field bindweed          Puncturevine          Tawha (morningglory) and many other perennial broadleaf weeds	Add 1/2 gallon of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	For effective broad spectrum control of annual and perennial broadleaf weeds and vines apply Brush Buster Herbicide when weeds are actively growing. Apply at the rate of 50 gallons of spray mix per 1/2 acre (about 22,000 square feet or equivalent to a strip 5,500 feet long x 4 feet wide). If less than 50 gallons of spray mix is applied per half acre, increase the concentration of Brush Buster Herbicide to maintain the same rate of active ingredients per half-acre.
<b>Annual Broadleaf Weeds</b> Pigweed                  Herbit                  Ragweed Chickweed              Knotweed              Sheep sorrel Clover                   Lambquarters          (red sorrel) Cocklebur              Mustard               Smartweed Dogfennel              Pigweed                Velvetleaf English daisy          Purslane               Wild buckwheat	Add 1 quart of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1/2 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	<b>FOR SPOT TREATMENTS</b> Mix 1 to 2 ounces of Brush Buster Herbicide with 2 to 3 gallons of water to treat 700 square feet.
<b>Woody Brush</b> Alder                      Hemlock                Serviceberry Ash                        Hornbeam              Spicebush Aspen                      Locust                   Spruce Basswood                Maple                    Sumac Cedar                      Mesquite               Sycamore Cherry                      Oak                       Thornapple Chinquapin              Persimmon             Thornberry Cucurbitaceae          Pine                      Wild plum Gum                        Poplar                   Willow Guava                      Sassafras              Witchhazel Dogwood                Schinus                 Yaupon Elm                        (Christmasberry) and similar woody plant species	<b>STEM FOLIAGE APPLICATION—HIGH WATER VOLUME</b>	
	Add 1/2 gallon of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost. Treat all stems and foliage to runoff with special emphasis on covering the root crown. Apply at the rate of 50-75 gallons of spray mix per 1/2 acre (about 11,000 square feet) depending on the height and density of the brush.
	<b>AERIAL APPLICATION* — 15 GALLONS OF SPRAY MIX/ACRE</b>	
	Add 3 gallons of Brush Buster Herbicide (equivalent to 3 lbs dicamba plus 6 lbs 2,4-D acid) plus 12 gallons of water to make 15 gallons of spray mix	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost. Apply at the rate of 15 gallons of spray mix per acre of brush.
<b>AERIAL APPLICATION* — 30 GALLONS OF SPRAY MIX/ACRE</b>		
Add 3 gallons of Brush Buster Herbicide (equivalent to 3 lbs dicamba plus 6 lbs 2,4-D acid) to 27 gallons of water to make 30 gallons of spray mix.	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost. Apply at the rate of 30 gallons of spray mix per acre of brush.	

\*Before making aerial applications, consult your State or Local Authorities for possible aerial application restrictions. To reduce the hazards of spray drift when applying this product by ground or air equipment, apply with a drift reducing agent such as foam spray adjuvant, inert system or with other systems designed to reduce spray drift.

BRUSH BUSTER IS A TRADEMARK OF VELSICOL CHEMICAL CORPORATION