

Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have stopped. Bait placements should be checked daily and replace consumed, contaminated or spoiled baits. The baits being replaced should immediately be disposed of in such a manner to prevent ingestion by children, pets, wildlife, and domestic animals.

Bait placement should be kept permanently to prevent new infestation.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in dry, cool area separately from fertilizer, feed, or foodstuffs and away from products with strong odors. Spillage should be carefully swept up and collected for disposal.

PESTICIDE DISPOSAL: Pesticide wastes may be toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse plastic pails. For paper or plastic bags, or for fiber drums with liners, completely empty into application equipment by shaking and tapping container. Offer pails or drums for recycling or reconditioning or puncture before disposal. Dispose of bags, liners, or punctured containers in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied for results due to misuse, improper handling, or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

Ramik is a Registered Trademark of Velsicol Chemical Corporation

Ramik[®]

Green

Bait Pack

Water Resistant, Weather Resistant

Rodenticide

For Control of Commercial

Mice & Rats

Indoor and Outdoor Use



ACTIVE INGREDIENT:

Diphacinone, Diphacoumatilol
Insecticide, 0.002

INERT INGREDIENTS

TOTAL 0.002

NET WEIGHT 1.000 LBS

NET WEIGHT 453.6 G

NET
CONTENTS
1 POUND

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
READ LABEL FOR ADDITIONAL CAUTIONS

VELSICOL CHEMICAL CORPORATION

341 East Ohio Street, Chicago, IL 60611

CAUTION

Keep out of the reach of children, pets, wildlife and domestic animals.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K₁, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

DIRECTIONS FOR USE

This product can be used in wet or damp locations such as river banks, railroad tracks, loading areas, in or around rat holes, under granaries, along fences, garbage dumps, etc.

This product is equally effective in dry areas such as the insides of buildings and other protected areas.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals or in tamperproof bait boxes.

Place baits along walls, in corners, in concealed locations, next to burrows or in runways where rats and mice feed, water, and travel.

For Norway and Roof Rats, place one to four bait packs per placement. Placements should be made at 15-foot intervals.

For House Mice, open the bait pack and place 1/4 to 1/2 ounce of bait at each placement. Placements should be made at 6- to 10-foot intervals.

Continued next column

876-208

05/01/1985

1/2

ACCEPTED

MAY 1 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. *076-208*