

VELSICOL[®] Vegatrol A-20-2T

HERBICIDE FOR BRUSH CONTROL

Contains 2 lbs. 2,4-D and 2 lbs. 2,4,5-T Acid Equivalents Per Gallon

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxy-
acetic Acid* 24.7%

Triethylamine Salt of 2,4,5-Trichlorophenoxy-
acetic Acid** 28.6%

INERT INGREDIENTS: 46.7%

TOTAL 100.0%

*Equivalent to 20.5% 2,4-Dichlorophenoxyacetic Acid.

**Equivalent to 20.5% 2,4,5-Trichlorophenoxyacetic Acid.

E.P.A. Reg. No. 876-214-AA

NET CONTENTS:

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Precautions

CAUTION

Harmful if swallowed. Avoid inhaling spray mist. This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin flush with plenty of water; for eyes flush with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Prevent drift of product to desirable plants, particularly beans, clover, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf crops during both the growing and dormant periods. Use coarse sprays which are less likely to drift.

Do not apply by airplane when sensitive crops are nearby. Consult your State or Local Authorities for possible aerial application or other application restrictions.

Do not use around the home, recreation sites or similar areas.

Do not contaminate irrigation ditches or water used for domestic purposes.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

To avoid injury to desirable plants, equipment used for Vegatrol A-2D-2T Herbicide should be thoroughly cleaned before reuse to handle or apply other agricultural chemicals.

Do not allow this product to come in direct contact with seeds, insecticides or fungicides. Do not contaminate feed or foodstuffs. Do not mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

CONTAINER DISPOSAL

Do not reuse empty container. Empty containers should be rinsed with water at least three times before disposal and rinse water should be added to the spray tank. Discard container in a safe place. Do not burn.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

VELSICOL
VEGATROL A-2D-2T
HERBICIDE FOR
WEED CONTROL

ACCEPTED

NOV 21 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 876-214

CAUTION

KEEP OUT OF REACH OF CHILDREN

VELSICOL CHEMICAL CORP.

Vegatrol A-2D-2T will kill or be susceptible to 2,4-D and 2,4,5


- Alder (
- Aspen (
- Birch (
- Blackberry (
- Box elder (
- Buckbrush (
- Cherokee rose (
- Choke cherry (
- Coffee bean (

Mix the recommended amount for good spray coverage. DO NOT spray when plants are actively growing. To assure good spray mix continuously.

Do not graze dairy animals
Do not graze meat animals

Area or Type of Treatment
<p style="font-weight: bold; font-size: 0.8em;">Foliage Treatment</p> <p style="font-size: 0.8em;">(Pasture and noncrop areas, rights-of-way (utility, highway, pipeline, railroad), fencerows, wasteland, pasture and similar noncropland.)</p>

Comments



VELSICOL Vegatrol A-2D-2T HERBICIDE FOR BRUSH CONTROL

CAUTION
KEEP OUT OF REACH OF CHILDREN

VELSICOL CHEMICAL CORP.

ACCEPTED
NOV 21 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 876-214

Vegatrol A-2D-2T will kill or control the following as well as many other noxious plants susceptible to 2,4-D and 2,4,5-T.

Alder	Cottonwood	Locust	Sagebrush
Aspen	Curly indigo	Mexican weed	Sassafras
Birch	Elder	Mulberry	Sumac
Blackberry	Elderberry	Oak	Sweet gum
Box elder	Elm	Osage orange	Wild cherry
Buckbrush	Hawthorn	Persimmon	Wild grape
Chicksee rose	Hickory	Poison ivy	Wild plum
Choke cherry	Honeysuckle	Poison oak	Willow
Coffee bean	Indigo	Poplar	

DIRECTIONS FOR USE

Mix the recommended amount of Vegatrol A-2D-2T Herbicide with sufficient water to obtain good spray coverage. **DO NOT DILUTE WITH OIL.**


Best results are obtained when applications are made while plants are well leaved and actively growing. To assure uniform results, shake container prior to opening and agitate spray mix continuously.

Do not graze dairy animals on treated areas within 6 weeks after application.

Do not graze meat animals on treated areas within 2 weeks of slaughter.

Area or Type of Treatment	Rate of Vegatrol A-2D-2T	Application Directions and Limitations
Foliage Treatment Pasture and noncrop areas, rights-of-way (utility, highway, pipeline, railroad), fencerows, wasteland, pasture and similar noncrop and.	Dilute 1 gallon (2 lbs. 2,4-D and 2 lbs. 2,4,5-T acid equivalent) with 99 gallons of water for ground application. For aerial application, dilute with 4-19 gallons of water.	For control of plants up to 6 to 8 feet in height. Foliage and bark should be thoroughly wet with a coarse spray. Repeat applications may be necessary if new growth occurs. Best results are obtained when applications are made while the plants are well leaved and actively growing.

Comments



ACCEPTED WITH COMMENTS