

CAUTION

Harmful if swallowed. Avoid inhaling spray mist. This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin flush with plenty of water; for eyes flush with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth. Vapor from this product may injure susceptible plants in the immediate vicinity.

Prevent drift of product to desirable plants, particularly beans, clover, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf crops during both the growing and dormant periods. Use coarse sprays which are less likely to drift.

Do not apply by airplane when sensitive crops are nearby. Consult your State or Local Authorities for possible aerial application or other application restrictions.

Do not contaminate irrigation ditches or water used for domestic purposes.

This product is toxic to fish. Keep out of any body of water not intended for aquatic weed control. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

To avoid injury to desirable plants, equipment used for Vegatrol BE-4D Herbicide should be thoroughly cleaned before reuse to handle or apply other agricultural chemicals.

Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate feed or foodstuffs. Do not mix with liquid fertilizers.


DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

CONTAINER DISPOSAL

Do not reuse empty container. Empty containers should be rinsed with water at least three times before disposal and rinse water should be added to the spray tank. Discard container in a safe place. Do not burn.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.




14137 460AB

VELSICOL Vegatrol BE-4D

HERBICIDE

VEGATROL BE-4D	60.7
INERT INGREDIENTS	39.3
TOTAL	100.0%



CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

341 E. OHIO STREET, CHICAGO, ILLINOIS 60611

Vegatrol BE-4D Herbicide will kill plants susceptible to 2,4-D.

Alligator weed	Canada thistle
Arrowweed	Catnip
Artichoke	Chickweed
Bindweed (hedge, field & European)	Chicory
Bitterweed	Cocklebur
Boxelder	Coffeebean
Buckhorn	Creeping
Bull thistle	Curly dandelion
Bulrush	Duckweed
Burdock	Elderberry
Bur ragweed	Goldenrod
Buttercup	Ground ivy
	Hemp

Mix the recommended amount of good spray coverage. Add the herbicide. Mix well; then add the water. Best results are obtained with growing. Perennial and other weeds give adequate control.

For ground application use in the following manner:
For aerial application use in the following manner:

Use
Rights-of way (utility, highway, pipeline, railroad), fence, wasteland and similar uncultivated land.
Lakes, ponds, drainage ditches, banks (aquatic weeds)
Ornamental Turf
Pastures and rangeland
Fall seeded wheat, oats, barley
Spring seeded wheat, oats, barley
Corn, Sorghum (post emergence)
Corn (pre emergence)

Comments

VELSICOL

Vegatrol BE-4D

HERBICIDE

ACTIVE INGREDIENT

Bromoxynil 2,4-D acid equivalent

Percent Acid

65.7

INERT INGREDIENTS

34.3

TOTAL

100.0

ACCEPTED

NOV 21 1967

U.S. DEPT. OF AGRICULTURE

WASH. D.C.

20250

876-217

NET CONTENTS

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

341 E. OHIO STREET CHICAGO, ILLINOIS 60601

Vegatro BE-4D Herbicide will kill or control the following in addition to many other noxious plants susceptible to 2,4-D.

Alligator weed	Canada thistle	Hoary cress	Pigweed	Stinkweed
Arrowweed	Catnip	Honeysuckle	Plantain	Sumac
Artichoke	Chickweed	Indigo	Poison ivy	Sunflower
Bindweed (hedge, field & European)	Chicory	Ironweed	Pokeweed	Virginia creeper
Bitter watercress	Cocklebur	Jimsonweed	Povertyweed	Waterhyacinth
Boxelder	Coffeebean	Lambsquarters	Puncturevine	Waterhily
Buckhorn	Creeping Jenny	Locoweed	Purslane	Waterprimrose
Bull thistle	Curly indigo	Mexicanweed	Rush	Wild garlic
Bulrush	Duckweed	Morning glory	Russian thistle	Wild lettuce
Burdock	Elderberry	Mustard	Sagebrush	Wild onion
Bur rag	Goldenrod	Nutgrass	Sheepspurse	Wild radish
Buttercup	Ground ivy	Parrotfeather	Smartweed	Willow
	Hemp	Pennywort	Sowthistle	

DIRECTIONS FOR USE

Mix the recommended amount of Vegatrol BE-4D Herbicide with sufficient water to obtain good spray coverage. Add the Vegatrol BE-4D to about one-half the volume of water to be used. Mix well; then add the rest of the water and mix until spray mixture is uniform.

Best results are obtained when spraying is done while the weeds are young and actively growing. Perennial and other hard-to-control weeds may require a repeat application to give adequate control.

For ground application use in 20 to 100 gallons of water per acre.

For aerial application use in 5 to 20 gallons of water per acre.

Use	Rate of Vegatrol BE-4D Per Acre	Application Directions and Limitations
Rights-of-way (utility, highway, pipeline, railroad), fence rows, wasteland and similar non-cropland.	1 to 4 pints (1/2 to 2 lbs. acid equivalent).	Do not graze dairy animals on treated areas within seven days after application. NOTE: Herbicide treatment of public waters requires a permit from the appropriate agencies in most states. Contact your State Conservation Department or Game and Fish Commission for aid in securing a permit if one is required in your state. Wet foliage thoroughly. Make application when leaves are fully developed above the water line and plants are actively growing. Do not apply to more than 1/2 to 1/2 of a lake or pond in any one month, because excessive decaying vegetation may deplete the oxygen content of water, killing fish. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. NOTE: Fish may be killed at application rates recommended on this label.
Lakes, ponds, drainage ditch banks (aquatic weeds)	2 1/2 to 4 1/2 pints (1 1/4 to 2 1/4 lbs. 2,4-D acid equivalent)	
Ornamental Turf	1 to 3 pints (1/2 to 1 1/4 lbs. 2,4-D acid equivalent) depending on type and stage of weed growth	Do not use on Dichondra, Bent, Carpetgrass, Bermuda, St. Augustine grass or clover lawns. Do not treat newly seeded turf until after the second mowing and the lower dosage rate should be used.
Pastures and rangeland	1 to 4 pints (1/2 to 2 lbs. 2,4-D acid equivalent) depending on type and stage of weed growth	Use only on established stands of perennial grasses. Do not graze dairy animals on treated areas within seven days after application.
Fall seeded wheat, oats and barley	1/2 to 3/4 pints (1/4 to 3/8 lbs. 2,4-D acid equivalent)	Apply in early spring when weeds are small and before crop has reached the boot stage. Do not forage or graze treated fields within 2 weeks after application. Do not feed treated straw to livestock.
Spring seeded wheat, oats and barley	1/2 pint (1/4 lb. 2,4-D acid equivalent)	Apply after the fully tillered stage, except during the boot to dough stage. Do not forage or graze treated fields within 2 weeks after application. Do not feed treated straw to livestock.
Corn, Sorghum (post emergence)	1/2 pint (1/4 lb. 2,4-D acid equivalent)	Make application when weeds are actively growing. Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall or sorghum is 4 to 12 inches tall. Do not apply from tasseling to dough stage. Use drop nozzles when either crop is over 10 inches tall.
Corn (pre emergence)	1 to 2 quarts (1 to 2 lbs. 2,4-D acid equivalent)	Use higher rates on heavy soils. Do not use on sandy soils. Plant corn as deep as feasible.

ACCEPTED WITH COMMENTS