

CAUTION

Harmful if swallowed. Avoid inhaling spray mist. This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin flush with plenty of water; for eyes flush with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth. Vapors from this product may injure susceptible plants in the immediate vicinity.

Prevent drift of product to desirable plants, particularly beans, clover, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf crops during both the growing and dormant periods. Use coarse sprays which are less likely to drift.

Do not apply by airplane when sensitive crops are nearby. Consult your State or Local Authorities for possible aerial application or other application restrictions.

Do not use around the home, recreation sites or similar areas.

Do not contaminate irrigation ditches or water used for domestic purposes.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

To avoid injury to desirable plants, equipment used for Vegatrol BE-4T Herbicide should be thoroughly cleaned before reuse to handle or apply other agricultural chemicals.

Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate feed or foodstuffs. Do not mix with liquid fertilizers.


DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

CONTAINER DISPOSAL

Do not reuse empty container. Empty containers should be rinsed with water at least three times before disposal and rinse water should be added to the spray tank. Discard container in a safe place. Do not burn.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.



14129-460AB

VELSICOL Vegatrol BE-4T

HERBICIDE FOR BRUSH CONTROL

Contains 4 lbs. 2,4,5-T Acid Equivalent Per Gallon

ACTIVE INGREDIENT

Butoxyethyl Ester of 2,4,5-Trichlorophenoxy

Acetic Acid 62.5%


INERT INGREDIENTS

37.5%

TOTAL 100.0%

Equivalent to 44.9% 2,4,5-Trichlorophenoxy Acid

NET CONTENTS



CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

111 E. COOK STREET, CHICAGO, ILLINOIS 60601

Vegatrol BE-4T Herbicide v
plants susceptible to 2,4,5

- Ash
- Aspen
- Birch
- Black oak
- Blackjack oak
- Buckbrush
- Bur oak
- Choke cherry
- Cottonwood

For use in noncrop areas
rows and wasteland.

Vegatrol BE-4T can be mixed
Number 2 S. B. (oil) or k

When water is used as a carrier
with one-half of the volume
the water and mix until spray
wet with a coarse spray. E
plants are well leaved and
growth appears.

For control of woody plants
preferred.

Do not graze dairy animals

Do not graze meat animals

Area or Type of Treatment
Basal Bark Treatment
Frill Treatment
Stump Treatment
Rangeland: Mesquite Treatment (Oil Water Aerial Treatment)
Rangeland: Hickory, Blackjack, Scrub oaks (Oil Water Aerial Treatment)

Comments

Vegatrol BE-4T Herbicide will kill or control the following in addition to many other noxious plants susceptible to 2,4,5-T.

Ash	Currant	Locust	Poplar
Aspen	Dogwood	Maple	Post oak
Birch	Elder	Mesquite	Prickly pear
Black oak	Elm	Osage orange	Red oak
Blackjack oak	Hackberry	Pecan	Scrub oak
Buckeye	Hawthorn	Persimmon	Sumac
Bur oak	Hickory	Pin oak	Sweet gum
Chokeberry	Ironwood	Poison ivy	Wild blackberry
Cottonwood	Larkspur	Poison oak	

DIRECTIONS FOR USE

For use in noncrop areas such as rights-of-way (utility, highway, pipeline, railroad), fence-rows and wasteland.

Vegatrol EE-4T can be mixed with either water or oils such as diesel oil, Number 1 or Number 2 S. B. (oil) or kerosene, or as an oil-water spray when required.

When water is used as a carrier or diluent, mix the recommended dosage of Vegatrol BE-4T with one half of the volume required for adequate coverage. Mix well, then add the rest of the water and mix until spray mixture is uniform. Foliage and bark should be thoroughly wet with a coarse spray. Best results are obtained when applications are made while the plants are well leafed and actively growing. Repeat applications may be necessary if new growth appears.

For control of woody plants over 8 feet in height, basal, stump, or frill treatment is usually preferred.

Do not graze dairy animals on treated areas within 6 weeks after application.

Do not graze meat animals on treated areas within 2 weeks of slaughter.

Area or Type of Treatment	Rate of Vegatrol BE-4T	Application Directions and Limitations
Basal Bark Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 4 gallons).	Apply using a knapsack sprayer to cover bark around the trunk and stems to a height of 12 to 15 inches above soil line. Cover bark thoroughly even to the point of runoff at the soil line. Application can be made any time of the year except when snow or water prevent application at the soil line.
Frill Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 4 gallons).	Application may be made during any season. For control of plants having trunks more than 6 inches in diameter, make overlapping cuts in the bark around the trunk as close to the ground as possible. Treat fresh cuts with diluted mixture.
Stump Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 3 gallons).	Apply to freshly cut stumps of large trees or woody plants. Make application with a knapsack sprayer having a nozzle of medium orifice or by brush or swab. For best results, both sides and top of stump should be thoroughly wet. Application can be made any time of the year except when snow or water prevents application at the soil line. Repeat applications if regrowth occurs.
Rangeland Mesquite Treatment (Oil Water Aerial Treatment)	Dilute 1 gallon (4 lbs. 2,4,5-T acid equivalent) with 5 gallons of oil and then mix with 18 gallons of water.	This emulsion should be applied at the rate of 4 gallons per acre (2/3 lb. 2,4,5-T acid equivalent per acre). Where both an overstory and understory of hardwoods are present, a second application may be necessary for control.
Rangeland Hickory, Blackjack, Scrub oaks (Oil Water Aerial Treatment)	Dilute 1 gallon (4 lbs. 2,4,5-T acid equivalent) with 2 gallons oil and then mix with 7 gallons of water.	This emulsion should be applied at the rate of 5 gallons per acre. (2 lbs. 2,4,5-T acid equivalent per acre.) Where both an overstory and understory of hardwoods are present, a second application may be necessary for control.

ACCEPTED WITH COMMENTS