



CAUTION

Harmful if swallowed. Avoid inhaling spray mist. This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin flush with plenty of water; for eyes flush with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Prevent drift of product to desirable plants, particularly beans, clover, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf crops during both the growing and dormant periods. Use coarse sprays which are less likely to drift. Vapors from this product may injure susceptible plants in the immediate vicinity.

Do not apply by airplane when sensitive crops are nearby. Consult your State or Local Authorities for possible aerial application or other application restrictions.

Do not contaminate irrigation ditches or water used for domestic purposes.

This product is toxic to fish. Keep out of any body of water not intended for aquatic weed control.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

To avoid injury to desirable plants, equipment used for Vegatrol LV-4TP Herbicide should be thoroughly cleaned before reuse to handle or apply other agricultural chemicals.

Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate feed or foodstuffs. Do not mix with liquid fertilizers.


DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

CONTAINER DISPOSAL

Do not reuse empty container. Empty containers should be rinsed with water at least three times before disposal and rinse water should be added to the spray tank. Discard container in a safe place. Do not burn.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.



14133-460AB

VELSICOL

Vegatrol LV-4TP

HERBICIDE

Contains 4 lbs. of Silvex Equivalent Per Gallon

ACTIVE INGREDIENT

Isooctyl Ester of Silvex 2 (2,4,5

Trichlorophenoxy) Propionic Acid

65.2%

INERT INGREDIENTS

34.8%

TOTAL

100.0%

Equivalent to 4 lbs. of Silvex 2 (2,4,5 Trichlorophenoxy) Propionic Acid

NET CONTENTS

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

341 E. ONTARIO STREET CHICAGO, ILLINOIS 60611

Vegatrol LV-4TP Herbicide w
plants susceptible to 2,4,5-T.

Blackjack oak
Ground ivy
Mesquite
Pin oak

Poison
Post
Red

Alligatorweed
Arrowhead

Broomweed
Buckhorn
Chicory
Cocklebur
Common chickweed

Cri
Cui
Cu
De

Mix the recommended amount of water to be used in spray thoroughly and continuously.

To assure uniform results, a mixes thoroughly.

Best results are obtained when actively growing. Repeat application use in

For ground application use in
For aerial application use in

USE	
Noncrop areas (rights-of-way, utility, highway, pipeline, railroad fence rows, wasteland)	ground applicat
	Poison i and wild blackba
	Spot treat—water

Use	Rat LV-
Lakes, ponds, drainage ditches (Aquatic weeds)	Dilute (4 to 100 gal)

Turf	Dilute (1 1/2 to 40 gal)
------	--------------------------

Rice	Dilute (1/4 to 1 gal)
------	-----------------------

Sugar Cane (Preemergence) (Louisiana Only) For control of barnyardgrass, crabgrass, and johnsongrass seedlings	Dilute 1 acid eq 15 gal per acre
--	----------------------------------

Sugar Cane (Preemergence) (Louisiana Only) broad leaved weed in stubble cane	Dilute 1 acid eq 15 gal per acre
--	----------------------------------

CERTIFIED

107

107

107

Common vetch

Vegatrol LV-4TP Herbicide will kill or control the following as well as many other noxious plants susceptible to 2,4,5-T.P

Woody Growth			
Blackjack oak	Poison ivy	Saltcedar	White oak
Ground ivy	Post oak	Sand shinnery oak	Wild blackberry
Mesquite	Red oak	Scrub oak	Yucca
Pin oak			
Aquatic Weeds			
Alligatorweed	Pickersweed	Spatterdock	
Arrowhead	Rush	Waterlily	
Broadleaved Weeds			
Broomweed	Croton weed	Lambsquarters	Pokeberry
Buckhorn	Curly dock	Mexicanweed	Ragweed
Chicory	Curly wildigo	Mouseear	Chickweed
Cocklebur	Dandelion	Plantain	Sunflower
Common chickweed			Wild lettuce

DIRECTIONS FOR USE

Mix the recommended amount of Vegatrol LV-4TP Herbicide to about one half the volume of water to be used in spraying. Mix well, then add the rest of the water and agitate thoroughly and continuously.

To assure uniform results, always shake the container prior to opening and agitate spray mixes thoroughly.

Best results are obtained when applications are made while the plants are well leafed and actively growing. Repeat application if regrowth occurs.

For ground application use in 20 to 100 gallons of water per acre.

For aerial application use in 5 to 20 gallons of water per acre.

USE		Rate of Vegetrol LV-4TP Per Acre	Application Directions and Limitations
Noncrop areas (rights-of-way, utility, highway, pipeline, railroad) fence rows, wasteland)	ground application	Dilute 3 to 4 quarts (3 to 4 lbs. acid equivalent) with 100 gallons of water.	For woody growth 6 to 8 feet in height. Cover foliage thoroughly with mixture. Do not graze dairy animals.
	Poison ivy and wild blackberry	Dilute 2 quarts (2 lbs. acid equivalent) with 100 gallons of water.	
	Spot Treat—water	Dilute 1/2 cup with 3 gallons of water.	

Use	Rate of Vegetrol LV-4TP per Acre	Application Directions and Limitations
Lakes, ponds, drainage ditches (Aquatic weeds)	Dilute 1 to 2 gallons (4 to 8 lbs. acid equivalent) with 50 to 100 gallons of water.	Wet stems and leaves thoroughly. Application should be made when leaves are fully developed about the water line and plants are actively growing. Do not apply to more than 1/3 to 1/2 of a lake or pond in any one month, because excessive decaying vegetation may deplete oxygen content of the water, killing fish. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. NOTE: Fish may be killed at application rates recommended on this label. Herbicide treatment of public waters requires a permit from the appropriate agencies in most states. Contact your State Conservation Department or Game and Fish Commission for aid in securing a permit if one is required in your state.
Turf	Dilute 1 1/2 to 2 quarts (1 1/2 to 2 lbs. acid equivalent) with 20 to 40 gallons of water.	Best weed control is obtained with spring and fall application and weeds are actively growing. Turf damage is more apt to result in hot midsummer. DO NOT use on Dichondra, Carpetgrass, Bent, Bermuda, St Augustine grass or on clover lawns. Newly seeded turf should not be treated until after the second mowing and at the lower dosage rate.
Rice	Dilute 1 to 3 pints (1/4 to 3/8 lbs. acid equivalent) per acre	Use sufficient water to give good coverage, depending on species, age and growth conditions. Delay spraying until the second or third week after flooding (7-9 weeks after seeding), because rice plants are sensitive to 2,4,5 TP in its early stages of growth. Water in the field should be shallow enough to permit direct application of the spray mixture to the weeds. Make all treatments well in advance of heading.
Sugar Cane (Preemergence) (Louisiana Only) For control of barnyardgrass, crabgrass, and Johnsongrass seedlings	Dilute 1 quart (1 lb. acid equivalent) with 15 gallons of water per acre	Apply over the row shortly before the seeds are expected to germinate. If cane is shaved and off-barred, treat immediately following this operation.
Sugar Cane (Preemergence) (Louisiana Only) broad leaved weed in stubble cane	Dilute 1 quart (1 lb. acid equivalent) with 15 gallons of water per acre	Apply over the row when weeds are actively growing but before they send up a seed stalk. A second application may be required.

ACCEPTED WITH COMMENT