




Commento



CAUTION

Harmful if swallowed. Avoid inhaling spray mist. This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin flush with plenty of water; for eyes flush with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

Do not apply (except as recommended) on or near desirable plants or in locations where the chemical may be washed or moved into contact with their roots. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth. Vapors from this product may injure susceptible plants in the immediate vicinity.

Prevent drift of product to desirable plants, particularly beans, clover, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf crops during both the growing and dormant periods. Use coarse sprays which are less likely to drift.

Do not apply by airplane when sensitive crops are nearby. Consult your State or Local Authorities for possible aerial application or other application restrictions.

Do not use around the home, recreation sites or similar areas.

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

To avoid injury to desirable plants, equipment used for Vegatrol LV-4T Herbicide should be thoroughly cleaned before reuse to handle or apply other agricultural chemicals.


Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate feed or foodstuffs. Do not mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

CONTAINER DISPOSAL

Do not reuse empty container. Empty containers should be rinsed with water at least three times before disposal and rinse water should be added to the spray tank. Discard container in a safe place. Do not burn.

NOTICE—Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.



14128-460AB

VELSICOL Vegatrol LV-4T

HERBICIDE FOR BRUSH CONTROL

Contains 4 lbs. 2,4,5-T Acid Equivalent Per Gallon

ACTIVE INGREDIENT

Isooctyl ester of 2,4,5-Trichlorophenoxy-
acetic acid 65.3%


INERT INGREDIENTS:

34.7%

TOTAL 100.0%

Equivalent to 4.547 lbs. 2,4,5-Trichlorophenoxyacetic Acid

NET CONTENTS



CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

341 E. OHIO STREET, CHICAGO, ILLINOIS 60611

Vegatrol LV-4T will kill or control susceptible to 2,4,5-T.

Ash	Currant
Aspen	Dogwood
Birch	Elder
Black oak	Elm
Black locust	Hackberry
Buckbrush	Hawthorn
Bur oak	Hickory
Choke cherry	Ironwood
Cottonwood	Larkspur

Vegatrol LV-4T can be mixed with Number 2 fuel oil or kerosene

When water is used as a carrier with one-half of the volume of the water and agitate thoroughly

When used as an oil-water spray separate container (not the sprayer) in contact with this mixture of water in the spray tank. Mix thoroughly spraying to assure a uniform

Vegatrol LV-4T may be used on areas along fences, rights-of-way and foliage thoroughly. Best results are obtained by growing. Repeat applications

Do not graze dairy animals on treated areas
Do not graze meat animals on treated areas

Area or Type of Treatment	Restrictions
Basal Bark Treatment	Do not apply to trees with diameter less than 4 inches
Frill Treatment	Do not apply to trees with diameter less than 4 inches
Stump Treatment	Do not apply to trees with diameter less than 4 inches
Forest and noncrop areas (Aerial or ground application)	Do not apply to trees with diameter less than 4 inches
Pasture and rangeland (ground or aerial application)	Do not apply to trees with diameter less than 4 inches

VELSICOL Vegatrol LV-4T

HERBICIDE FOR BRUSH CONTROL

Contains 4 lbs. 2,4,5-T Acid Equivalent Per Gallon

ACTIVE INGREDIENT

Isooctyl ester of 2,4,5-Trichlorophenoxy-
acetic acid 65.3%

INERT INGREDIENTS: 34.7%

TOTAL 100.0%

Contains 4 lbs. 2,4,5-T Acid Equivalent Per Gallon

NET CONTENTS


CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel for Additional Cautions

VELSICOL CHEMICAL CORP.

341 E. OHIO STREET, CHICAGO, ILLINOIS 60611



Vegatrol LV-4T will kill or control the following as well as many other noxious plants susceptible to 2,4,5-T.

Ash	Currant	Locust	Poplar
Aspen	Dogwood	Maple	Post oak
Birch	Elder	Mesquite	Prickly pear
Black oak	Elm	Osage orange	Red oak
Black rock oak	Hackberry	Pecan	Scrub oak
Buckbrush	Hawthorn	Persimmon	Sumac
Bur oak	Hickory	Pin oak	Sweet gum
Chokecherry	Ironwood	Poison ivy	Wild blackberry
Cottonwood	Larkspur	Poison oak	

DIRECTIONS FOR USE

Vegatrol LV-4T can be mixed with either water or oils such as diesel oil, Number 1 or Number 2 fuel oil or kerosene, or as an oil-water spray when required.

When water is used as a carrier or diluent, mix the recommended dosage of Vegatrol LV-4T with one-half of the volume required for adequate coverage. Mix well, then add the rest of the water and agitate thoroughly and continuously.

When used as an oil-water spray, add the required amount of Vegatrol LV-4T to the oil in a separate container (not the spray tank). Do not allow any water or oil-water mixture to come in contact with this mixture as it may gel. Add this mixture to half the required volume of water in the spray tank. Mix well, then add the remaining water. Continue mixing while spraying to assure a uniform emulsion.

Vegatrol LV-4T may be used to control trees and brush in forest management, noncrop areas (along fences, rights-of-way and similar areas) pastures and rangelands. Wet bark and foliage thoroughly. Best results are obtained when plants are well leafed and actively growing. Repeat applications may be necessary if new growth occurs.

Do not graze dairy animals on treated areas within 6 weeks after application.

Do not graze meat animals on treated areas within 2 weeks of slaughter.

Area or Type of Treatment	Rate of Vegatrol LV-4T	Application Directions and Limitations
Basal Bark Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 4 gallons).	Apply using a knapsack sprayer to cover bark around trunk and stems from soil surface to a height of 12 to 15 inches. Cover bark thoroughly even to the point of runoff at the soil line. Application can be made any time of the year except when snow or water prevents application at the soil line.
Frill Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 4 gallons).	Application may be made during any season. For control of plants having trunks more than 6 inches in diameter, make overlapping cuts in the bark around the trunk as close to the ground as is possible. Treat fresh cuts with the diluted mixture.
Stump Treatment	Dilute 3 gallons (12 lbs. 2,4,5-T acid equivalent) with 97 gallons of oil (equivalent to 1 pint in 4 gallons).	Apply to freshly cut stumps of large trees or woody plants. Make application with a knapsack sprayer having a nozzle of medium orifice or by a brush or swab. For best results, both sides and top of stump should be thoroughly wet. Application can be made any time of the year except when snow or water prevents application at the soil line. Repeat applications if regrowth occurs.
Forest and noncrop areas (Aerial or ground application)	4 pints (2 lbs. 2,4,5-T acid equivalent) per acre.	Use sufficient volume of diluent to obtain good coverage. For ground application use in 20 to 100 gallons of diluent per acre. For aerial application, use 5-20 gallons of diluent per acre.
Pasture and rangeland (ground or aerial application)	4 pints (2 lbs. 2,4,5-T acid equivalent) per acre.	For ground application, use in 20 to 100 gallons of diluent per acre. For aerial application, use 5-20 gallons of diluent per acre. Do not apply to newly seeded areas. Observe grazing restrictions given above.

ACCEPTED WITH COMMENTS