

CAUTION

Keep away from children, pets, wildlife, and domestic animals. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling, and conditions of use, which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied, for results due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling, or storage of this material.

VELSICOL[®]

Ramik[®] Brown

APPLE FLAVORED RODENTICIDE PELLETS FOR REPACKAGING OF A RODENTICIDE

ACTIVE INGREDIENT:

Diphacinone (2-Diphenylacetyl-1, 3-Indandione)

0.005%

INERT INGREDIENTS:

99.995%

TOTAL 100.000%

E.P.A. Reg. No. 876-240-AA


NET CONTENTS:

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See left panel for additional cautions.


VELSICOL CHEMICAL CORPORATION
341 East Ohio Street • Chicago, Illinois 60611



DIRECTIONS FOR USE

Directions for use for the repackaged product may be found on Velsicol Specimen Label Number SP-138.

NOTE: Additional data may be required to support registration of your product, if your labeling text or graphics are altered from the example on the specimen label, where the hazards or efficacy are affected by such alteration.



CAUTION

Keep out of reach of children, pets, wildlife and domestic animals.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

DISCLAIMER STATEMENT

BRAND NAME

APPLE FLAVORED
RODENTICIDE
PELLETS

Manufactured for:

DIRECTIONS FOR USE

This product can be used in wet or damp locations such as river banks, railroad tracks, loading areas, in or around rat holes, under granaries, along fences, garbage dumps, etc.

This product is equally effective in dry areas such as the insides of buildings and other protected areas.

All outdoor placements must be made in tamperproof bait boxes.

Place baits in tamperproof bait boxes along walls, in corners and in concealed locations where rats can feed undisturbed.

Bait placements should be checked daily to make sure that an uninterrupted supply of bait is available. Maintain an uninterrupted supply of bait for at least 15 days. Continue baiting until all signs of feeding have stopped.

NORWAY AND ROOF RATS: Place baits 20 to 30 feet apart. Use 4 – 16 ounces of bait per location.

HOUSE MICE: Place baits 8 – 10 feet apart. Use tablespoon amounts (1/4 to 1/2 ounce) per location.

Where a continuous source of infestation is present (eg. nearby dumps) permanent bait stations should be established and the bait replenished as needed.

Rats and mice must feed repeatedly on the bait for several days before death occurs.

Warning or directions for use, term, etc. described herein are not intended to constitute registration of this product.

Date 5/9/15

CAUTION

Keep out of reach of children, pets, wildlife and domestic animals.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

DISCLAIMER STATEMENT

BRAND NAME

APPLE FLAVORED
RODENTICIDE
PELLETS

Manufactured for:

DIRECTIONS FOR USE

This product can be used in wet or damp locations such as river banks, railroad tracks, loading areas, in or around rat holes, under granaries, along fences, garbage dumps, etc.

This product is equally effective in dry areas such as the insides of buildings and other protected areas.

All outdoor placements must be made in tamperproof bait boxes.

Place baits in tamperproof bait boxes along walls, in corners and in concealed locations where rats can feed undisturbed.

Bait placements should be checked daily to make sure that an uninterrupted supply of bait is available. Maintain an uninterrupted supply of bait for at least 15 days. Continue baiting until all signs of feeding have stopped.

NORWAY AND ROOF RATS: Place baits 20 to 30 feet apart. Use 4 – 16 ounces of bait per location.

HOUSE MICE: Place baits 8 – 10 feet apart. Use tablespoon amounts (1/4 to 1/2 ounce) per location.

Where a continuous source of infestation is present (eg. nearby dumps) permanent bait stations should be established and the bait replenished as needed.

Rats and mice must feed repeatedly on the bait for several days before death occurs.

Date