

CAUTION

Keep out of the reach of children, pets, wildlife and domestic animals. Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bis hydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN

IMPORTANT

For best results, keep this product in a closed container free from odors which may contaminate the bait and reduce acceptability. Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy when empty.

DIRECTIONS FOR USE

This product can be used in wet or damp locations such as river banks, railroad tracks, loading areas, in or around rat holes, under granaries, along fences, garbage dumps, etc.

This product is equally effective in dry areas such as the inside of buildings and other protected areas.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes.

Place baits along walls, in corners, in concealed locations, next to burrows or in runways where rats and mice feed, water and travel.

For Norway and Roof Rats, place one to four bait packs per placement. Placements should be made at 15-foot intervals.

For House Mice open the bait pack and place 1/4 to 1/2 ounce of bait at each placement. Placements should be made at 6- to 10-foot intervals.

Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have stopped. Bait placements should be checked daily and replace consumed, contaminated or spoiled baits. The baits being replaced should immediately be disposed of in such a manner to prevent ingestion by children, pets, wildlife and domestic animals.

Bait placement should be kept permanently to prevent new infestation.

NOTICE: Because Velsicol Chemical Corporation has no control over storage, handling and conditions of use which are of critical importance, Velsicol Chemical Corporation makes no representation or warranty, either express or implied for results due to misuse, improper handling or storage of this material. Nor does Velsicol Chemical Corporation assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

VELSICOL CHEMICAL CORPORATION

16-110-1006

341 East Ohio Street • Chicago, Illinois 60611

VELSICOL™

Ramik™ Brown*

Bait Packs

Apple Flavored Weather-Resistant Rodenticide
For Control of Commensal Mice and Rats
Indoors and Outdoors

ACTIVE INGREDIENT:

Diphacinone (2-Diphenylacetyl-1, 3
Indandione) 0.005%

INERT INGREDIENTS: 99.995%

TOTAL 100.000%

E.P.A. Reg. No. 876-247-AA

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See back panel for additional cautions.

Net Contents

4 ounces



ACCEPTED

MAY 13 1975

Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, for the pesticide registered under EPA Reg. No. 876-247

VELSICOL CHEMICAL CORPORATION

341 East Ohio Street • Chicago, Illinois 60611

CAUTION

Keep out of the reach of children, pets, wildlife and domestic animals.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K, combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdoses of bis hydroxycoumarin (Dicumarol).

IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

IMPORTANT

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce its acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

NOTICE: This product is a rodenticide. It is intended for use only against rodents. It is not intended for use against insects, birds, or other animals. It is not intended for use in areas where it may be ingested by children, pets, wildlife, or domestic animals. It is not intended for use in areas where it may be ingested by livestock. It is not intended for use in areas where it may be ingested by wild birds. It is not intended for use in areas where it may be ingested by wild mammals. It is not intended for use in areas where it may be ingested by wild reptiles. It is not intended for use in areas where it may be ingested by wild amphibians. It is not intended for use in areas where it may be ingested by wild fish. It is not intended for use in areas where it may be ingested by wild aquatic invertebrates. It is not intended for use in areas where it may be ingested by wild plants. It is not intended for use in areas where it may be ingested by wild fungi. It is not intended for use in areas where it may be ingested by wild bacteria. It is not intended for use in areas where it may be ingested by wild viruses. It is not intended for use in areas where it may be ingested by wild protozoa. It is not intended for use in areas where it may be ingested by wild algae. It is not intended for use in areas where it may be ingested by wild lichens. It is not intended for use in areas where it may be ingested by wild mosses. It is not intended for use in areas where it may be ingested by wild ferns. It is not intended for use in areas where it may be ingested by wild gymnosperms. It is not intended for use in areas where it may be ingested by wild angiosperms. It is not intended for use in areas where it may be ingested by wild fungi. It is not intended for use in areas where it may be ingested by wild bacteria. It is not intended for use in areas where it may be ingested by wild viruses. It is not intended for use in areas where it may be ingested by wild protozoa. It is not intended for use in areas where it may be ingested by wild algae. It is not intended for use in areas where it may be ingested by wild lichens. It is not intended for use in areas where it may be ingested by wild mosses. It is not intended for use in areas where it may be ingested by wild ferns. It is not intended for use in areas where it may be ingested by wild gymnosperms. It is not intended for use in areas where it may be ingested by wild angiosperms.

VELSICOL™

Ramik™ Brown*

Bait Packs

**Apple Flavored Weather-Resistant Rodenticide
For Control of Commensal Mice and Rats
Indoors and Outdoors**

ACTIVE INGREDIENT:

Diphacinone (2-Diphenylacetyl 1, 3-Indandione)

0.005%

INERT INGREDIENTS:

99.995%

TOTAL

100.000%


E.P.A. Reg. No. 876-247-AA

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See left panel for additional cautions.

Net Contents



VELSICOL CHEMICAL CORPORATION

341 East Ohio Street • Chicago, Illinois 60611

DIRECTIONS FOR USE

This product may be used in wet or damp locations such as river banks, railroad tracks, loading areas, in or around rat holes, under granaries, along fences, garbage dumps, etc.

This product is equally effective in dry areas such as the insides of buildings and other protected areas.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof bait boxes.

Place baits along walls, in corners, in concealed locations, next to burrows or in runways where rats and mice feed, water and travel.

For Norway and Roof Rats, place one to four bait packs per placement. Placements should be made at 15-foot intervals.

For House Mice open the bait pack and place 1/4 to 1/2 ounce of bait at each placement. Placements should be made at 6- to 10 foot intervals.

Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have stopped. Bait placements should be checked daily and replace consumed, contaminated or spoiled baits. The baits being replaced should immediately be disposed of in such a manner to prevent ingestion by children, pets, wildlife and domestic animals.

Bait placements should be kept permanently to prevent new infestation.